

Non-coding RNAs and resistance to anticancer drugs in gastrointestinal tumours

Jens C. Hahne¹, Nicola Valeri^{1, 2*}

¹Institute of Cancer Research (ICR), United Kingdom, ²Department of Medicine, Royal Marsden NHS Foundation Trust, United Kingdom

Submitted to Journal:
Frontiers in Oncology

Specialty Section:
Cancer Molecular Targets and Therapeutics

Article type:
Review Article

Manuscript ID:
376456

Received on:
20 Mar 2018

Revised on:
23 May 2018

Frontiers website link:
www.frontiersin.org

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Author contribution statement

NV and JH idea, conception and writing parts of the review.

Keywords

non-coding RNA, lncRNA, microRNA, anticancer drugs, gastrointestinal tumour, cancer therapy, Resistance

Abstract

Word count: 135

Non-coding RNAs are important regulators of gene expression and transcription. It is well established that impaired non-coding RNA expression especially the one of long non-coding RNAs and microRNAs is involved in a number of pathological conditions including cancer. Non-coding RNAs are responsible for the development of resistance to anticancer treatments as they regulate drug resistance-related genes, affect intracellular drug concentrations, induce alternative signalling pathways, alter drug efficiency via blocking cell cycle regulation and DNA damage response. Furthermore, they can prevent therapeutic-induced cell death and promote epithelial-mesenchymal transition and elicit non-cell autonomous mechanisms of resistance. In this review we summarise the role of non-coding RNAs for different mechanisms resulting in drug resistance (e.g. drug transport, drug metabolism, cell cycle regulation, regulation of apoptotic pathways, cancer stem cells and epithelial-mesenchymal transition) in the context of gastrointestinal cancers.

Funding statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Non-coding RNAs and resistance to anticancer drugs in gastrointestinal tumours

Jens C. Hahne¹, Nicola Valeri^{1,2,*}

¹ Division of Molecular Pathology, The Institute of Cancer Research London & Sutton, UK

² Department of Medicine, The Royal Marsden NHS Trust, London & Sutton, UK.

Correspondence:

Dr. Nicola Valeri

Centre for Molecular Pathology

The Institute of Cancer Research & The Royal Marsden NHS Foundation Trust

Cotswold Road, Sutton, Surrey, SM2 5NG, UK

Telephone: +44 0208 915 6634

Email: nicola.valeri@icr.ac.uk

Keywords: non-coding RNA, lncRNA, microRNA, anticancer drugs, gastrointestinal
tumour, cancer therapy, resistance

Word counts: 9117 – this article is written in British English

7 Figures

3 Tables

Abstract:

Non-coding RNAs are important regulators of gene expression and transcription. It is well established that impaired non-coding RNA expression especially the one of long non-coding RNAs and microRNAs is involved in a number of pathological conditions including cancer. Non-coding RNAs are responsible for the development of resistance to anticancer treatments as they regulate drug resistance-related genes, affect intracellular drug concentrations, induce alternative signalling pathways, alter drug efficiency via blocking cell cycle regulation and DNA damage response. Furthermore, they can prevent therapeutic-induced cell death and promote epithelial-mesenchymal transition and elicit non-cell autonomous mechanisms of resistance.

In this review we summarise the role of non-coding RNAs for different mechanisms resulting in drug resistance (*e.g.* drug transport, drug metabolism, cell cycle regulation, regulation of apoptotic pathways, cancer stem cells and epithelial-mesenchymal transition) in the context of gastrointestinal cancers.

Introduction:

Gastrointestinal (GI) cancer encompasses a heterogeneous group of tumours that affect the digestive tract system (Pourhoseingholi et al., 2015). These include cancers of the oesophagus, stomach, gallbladder, liver and biliary tract, pancreas, small intestine, colon, rectum and anus. GI cancer is the most common form of cancer responsible for nearly 25% of all new cancer diagnosis and responsible for most of cancer related death (around 30% of all cancer related death) worldwide (Siegel et al., 2015; Torre et al., 2015).

Chemotherapy is, alongside with surgery and radiation therapy, one of the main treatments for cancer (Hung et al., 2006; Chan et al., 2016; Ismael et al., 2016; Jakhetiya et al., 2016; Murphy, 2016; Olcina and Giaccia, 2016; Rautio et al., 2016; Ristamaki and Algars, 2016; Rutkowski and Hompes, 2016). Many chemotherapeutic agents have successfully prolonged overall and progression-free survival of GI cancer patients (Slamon et al., 2001; Motzer et al., 2007; Blanke et al., 2008; Maemondo et al., 2010; Chapman et al., 2011). In addition, a better understanding of the biology and mechanism underpinning GI cancer initiation and progression is leading to more personalised treatments. Indeed, identification of

well-defined molecular subtypes and/or molecular profiling of somatic mutations offer the opportunity to further optimize the efficacy of treatments through tailored approaches (Kwak et al., 2010; Douillard et al., 2013; Korpanty et al., 2014; Siroy et al., 2015).

Despite major improvements in the management of GI cancer patients, resistance to therapies arises almost inevitably at some point during the treatment and chemo-resistance is one of the main challenges in cancer therapy (Housman et al., 2014). Drug resistance can be caused by gene mutations, abnormal DNA repair, alteration in cell cycle regulation, cell death inhibition (mostly caused by deregulated apoptotic signalling pathways), reduced drug efficacy as well as enhanced drug clearance (Zahreddine and Borden, 2013; Housman et al., 2014). Furthermore, the epithelial-mesenchymal transition (EMT) process and the presence of tumour stem cells have been identified as causes of drug resistance (Shang et al., 2013; Xia and Hui, 2014; Mitra et al., 2015; Prieto-Vila et al., 2017). The complex molecular mechanisms of chemo-resistance have not been fully elucidated yet and a better understanding of drivers of primary and secondary resistance to chemotherapy will likely result into improved patients' survival. Increasing evidence points towards the role of non-coding RNAs as a central hub for treatment resistance. Therefore, this review outlines the role of non-coding RNAs for the different drug resistance mechanisms involved in GI cancer therapy failure. Table 1 summarised the non-coding RNAs discussed in this review and in figure 1-6 the role for each of these non-coding RNAs in the context of the different GI tumours is illustrated.

Non-coding RNAs:

In human tissues the amount of non-coding RNAs is more than three times higher compared to the amount of protein-coding RNAs (Geisler and Coller, 2013). Non-coding RNAs are a large family that includes more than 16 categories of long and short RNA molecules (Table 2); among them transfer RNAs (tRNAs), ribosomal RNAs (rRNAs), small nucleolar RNAs (snoRNAs), endogenous small interfering RNAs (endo-siRNAs), sno-derived RNAs (sdRNAs), transcription initiation RNAs (tiRNAs), miRNA-offset-RNAs (moRNAs), circular RNAs (circRNAs), vault RNAs (vRNAs), microRNAs, small interfering RNAs (siRNAs), small nuclear RNAs (snRNAs), extracellular RNAs (exRNAs), piwi-interacting RNAs (piRNAs), small Cajal body RNAs (scaRNAs), long intergenic non-coding RNAs

(lincRNAs) and long non-coding RNAs (lncRNAs), all of which are not coding for known proteins (Taal et al., 1993;Eddy, 2001;He and Hannon, 2004;Guttman et al., 2009;Langenberger et al., 2009;Taft et al., 2009a;Taft et al., 2009b;Wilusz et al., 2009;Choudhuri, 2010;Ling et al., 2013;Claycomb, 2014;Guo et al., 2014;An et al., 2016;Azlan et al., 2016;Beermann et al., 2016;de Almeida et al., 2016;Evans et al., 2016;Geiger and Dalgaard, 2016;Granados-Riveron and Aquino-Jarquin, 2016;Khurana et al., 2016;Qi et al., 2016;Quinn and Chang, 2016).

Long non-coding RNAs (lncRNAs) and microRNAs are the most studied non-coding RNAs playing a role in anticancer drug resistance and will be covered in this review.

LncRNAs are composed of more than 200 nucleotides. They are important regulators during development and pathological processes (Guttman et al., 2011;Sauvageau et al., 2013;Herriges et al., 2014;Li et al., 2014a;Ounzain et al., 2014). LncRNAs are pivotal in regulating gene expression by binding to chromatin regulatory proteins and they are able to alter chromatin modification as well as transcriptional or post-transcriptional gene regulation by interacting with other RNAs and proteins (Moran et al., 2012;Kornienko et al., 2013;Han and Chang, 2015). Recently, a crosstalk and strong linkage between lncRNA and microRNAs has been identified (Yoon et al., 2014). It has been shown that lncRNA stability can be reduced by interaction with specific microRNAs and, *vice versa*, lncRNAs act as microRNA decoys sequestering microRNAs from the intra-cellular cytosol and leading to re-expression of microRNA target genes (Yoon et al., 2014). Furthermore, lncRNAs can promote gene expression by competing with microRNAs for specific binding sites in the non-coding regions of mRNAs and prevent the transcriptional repression caused by microRNAs (Yoon et al., 2014). Interestingly some lncRNAs can be processed into microRNAs (Yoon et al., 2014) suggesting a plastic interaction among different classes of non-coding RNAs.

MicroRNAs are short RNA transcripts of 18–24 nucleotides. They are responsible for fine tuning cell homeostasis by controlling gene expression at post-transcriptional level, (Fabbri et al., 2009;Valeri et al., 2009;Winter et al., 2009). Due to the fact that each microRNAs can have several target mRNAs the interaction of one microRNA with various target mRNAs results in direct deregulation of different target proteins acting simultaneously in regulation of diverse cellular pathways (Macfarlane and Murphy, 2010;Pasquinelli, 2012). Therefore, variation in microRNA expression can result in reduced mRNA levels ultimately resulting in

changes in protein levels within the cell (von Schack et al., 2011;Pasquinelli, 2012). MicroRNAs expression patterns are tissue-specific (Lagos-Quintana et al., 2002) and often define the physiological status of the cell (Lim et al., 2005). Strong clinical and pre-clinical evidence suggests that microRNA aberrant expression plays a role in several diseases including cancer, infectious, neurodegenerative and immune-related diseases. (Murakami et al., 2006;Mitchell et al., 2008;O'Connell et al., 2010;Esteller, 2011;Ha, 2011b;a;c;Grasedieck et al., 2012;Iorio and Croce, 2012;Acunzo et al., 2015;Balatti et al., 2015;Gardiner et al., 2015). Analysis of microRNA expression patterns represents a promising tool for cancer diagnosis, prognosis and treatment prediction. MicroRNAs have been extensively studied in monitoring treatment resistance in consideration of their high stability in tissues and body fluids. In blood, microRNAs are included in RNA-binding multiprotein complexes and/or exosomes and their short length makes microRNAs less prone to degradation and improves their stability under different sample storage conditions in blood (Mitchell et al., 2008;Macfarlane and Murphy, 2010;Grasedieck et al., 2012;Gardiner et al., 2015) .

General principles of drug resistance:

Drug resistance is classified into intrinsic and acquired. Primary drug resistance is pre-existing and renders cancer cells immune against the therapy from the very beginning. In contrast, acquired (secondary) drug resistance develops during therapy due to adaptive processes of the tumour (Gottesman et al., 2002;Longley and Johnston, 2005;Rodrigues et al., 2012a;Holohan et al., 2013;Housman et al., 2014). Different mechanisms are involved in primary and acquired drug resistance and relate to non-coding RNAs dysregulation.

Deregulation of proteins involved in drug metabolism

One reason for drug resistance can be found on the level of drug transport. Reduced influx or increased efflux of chemotherapeutics result in lower intracellular drug concentrations and promotes therapy failure (Gottesman et al., 2002). Altered drug metabolism is another possible cause for drug resistance. Drug metabolism is a complex pathway composed of

multiple proteins for detoxification of foreign compounds (*e.g.* chemotherapeutics) normally neither produced nor present in a cell (Michael and Doherty, 2005). This pathway can be subdivided into modification (phase I reaction), conjugation (phase II reaction) and excretion (phase III reaction) (Park, 2001). Several drug-metabolizing enzymes, especially members of the cytochrome P450 family, together with drug transporters increase the polarity of the drugs during phase I (Shimada et al., 1989;Guengerich and Shimada, 1991). In the following phase II the polarity of the drugs is further increased by conjugation reactions (Shea et al., 1988b;McLellan and Wolf, 1999b). Finally, in phase III the resulting drug metabolites are exported by transmembrane transporter like ATP-binding cassette (ABC) proteins and solute carrier (SLC) transport proteins (Dean et al., 2001;Kathawala et al., 2015;Lin et al., 2015;Colas et al., 2016).

The vaults are known to contribute to drug resistance by transporting drugs away from their intracellular targets and vaults are involved in drug sequestration (Mossink et al., 2003). The vRNAs hvg-1 and hvg-2 that are present in the vaults (Table 2) interact with drugs via specific binding sites (Gopinath et al., 2010). In agreement with their role in regard to drug resistance the number of vaults are increased in cancer patients who developed resistance under chemotherapy (Mossink et al., 2003). In addition, the vRNAs are producing several small RNAs among them is svRNAb which down-regulates the key enzyme in drug metabolism CYP3A4 and accounts so for multidrug resistance in GI cancers (Persson et al., 2009).

Furthermore, lncRNA H19 was identified as another non-coding RNA involved in drug resistance. The oncogenic potential of lncRNA H19 was demonstrated in different tumour types (*e.g.* liver and oesophageal cancer) and over-expression of lncRNA H19 was observed in parallel with up-regulation of the membrane glycoprotein p95 in multidrug-resistant tumours (Tsang and Kwok, 2007;Matouk et al., 2013). In liver tumour cells resistant to doxorubicin, etoposide, paclitaxel and vincristine lncRNA H19 expression was increased (Tsang and Kwok, 2007). LncRNA H19 participate in the regulation of *MDR1* gene (also known as *ABCB1* gene) expression and modulate the drug transport out of the cell (Tsang and Kwok, 2007). *In-vitro* models of hepatocellular carcinoma suggest that lncRNA H19 can alter

MDR1 promoter methylation and, in doing so, increases the transcription of P-glycoprotein (Tsang and Kwok, 2007).

Similarly, in gastric cancer, lncRNA MRUL (MDR-related and up-regulated lncRNA) acts as an enhancer for transcription of P-glycoprotein (MDR1) (Wang et al., 2014) increasing the number of transmembrane transporters on the tumour cell membrane and fosters the drug export (Wang et al., 2014). As we described above, different non-coding RNAs can merge onto the same pathway: this is the case of lncRNA AK022798 whose expression is induced by NOTCH-1 over-expression during gastric cancer progression (Hang et al., 2015). LncRNA AK022798 in turn up-regulates the expression of P-glycoprotein and is responsible for increased cisplatin resistance in gastric cancer patients (Hang et al., 2015). Similarly, in cisplatin and 5-fluorouracil resistant gastric cancer patients the expression of lncRNA PVT-1 (plasmacytoma variant translocation 1) and lncRNA ANRIL (antisense to CDKN2B locus) are also increased and these non-coding RNAs promote MDR1 up-regulation and drug resistance (Zhang et al., 2015b; Lan et al., 2016).

Non-coding RNA dysregulation is tissue specific, indeed Wnt- β -catenin pathway activation triggers the expression of a different lncRNA, CCAL (colorectal cancer-associated lncRNA). The effect on phenotype is the same as in other cancers given CCAL in turn up-regulates P-glycoprotein expression and causing chemotherapy resistance (Ma et al., 2016b).

Additional to the regulation via lncRNAs ABC transporter expression levels are also controlled by miRNAs (Haenisch et al., 2014; Ikemura et al., 2014).

In colon cancer, P-glycoprotein expression was found to be directly deregulated at post-transcriptional level by binding of miR-145 to the 3'-UTR of the *MDR1* gene transcript (Ikemura et al., 2013). Down-regulation of miR-145 results in increased ABCB1 protein level (Ikemura et al., 2013). Analogously miR-297 binds to the 3'-UTR of ABCC2 mRNA and suppresses the expression of ABCC2 transporter (Xu et al., 2012). In chemo-resistant colorectal carcinoma, miR-297 is often down-regulated and consequently ABCC2 is expressed on a higher level compared to the surrounding colon tissue (Xu et al., 2012). Interestingly, *in-vitro* and *in-vivo* models suggest that resistance to vincristine and oxaliplatin could be overcome by restoring miR-297 expression in therapy resistant cells (Xu et al., 2012). Virtually expression of all the transporters can be affected by microRNA dysregulation; ABCB5 transporter is highly expressed in colon cancer cell lines with down-

regulated miR-522 expression and renders these cells resistant to doxorubicin treatment (Yang et al., 2015). MiR-522 binds to the ABCB5 mRNA 3'-UTR and over-expression of miR-522 reverse chemo-resistance to doxorubicin (Yang et al., 2015). Similarly, 5-fluorouracil resistance in microsatellite instable colon cancer (caused by deregulated miR-21 or miR-155 (Valeri et al., 2010a; Valeri et al., 2010b) as mentioned in detail later) can be enhanced by down-regulation of miR-23a resulting in higher expression of the direct target ABCF1 (Li et al., 2015d).

Similar examples exist across the board: in gastric cancer for example, down-regulation of miR-508-5p was identified as a reason for multidrug resistance (Shang et al., 2014). MiR-508-5p represses the expression of P-glycoprotein and the transcription factor zinc ribbon domain-containing 1 (ZNRD1) that is an important factor for *MDR1* gene translation (Shang et al., 2014). Loss of miR-508-5p decreased drug sensitivity in gastric cancer *in-vitro* and *in-vivo*, whereas ectopic expression of miR-508-5p overcomes drug resistance (Shang et al., 2014).

In pancreatic cancer cell lines, expression of the transporter ABCC1 is controlled by miR-1291 binding to the 3'-UTR (Pan et al., 2013). MiR-1291 is often down-regulated in pancreatic cancer resulting in an increased expression of ABCC1 that finally leads to higher efflux rate of toxic substances (Munoz et al., 2007; Tu et al., 2016). This is the reason for resistance to many chemotherapeutics, such as anthracyclines (*e.g.*, doxorubicin), platinum derivatives and the folate antagonist methotrexate (Munoz et al., 2007; Tu et al., 2016). Another transporter, called ATP7A (ATPase Cu²⁺ transporting alpha polypeptide), is up-regulated in *in-vitro* models of resistant pancreatic tumours due to decreased expression of miR-374b (Schreiber et al., 2016) and increased ATP7A protein expression is at least partially responsible for cisplatin resistance in pancreatic cancer model systems (Schreiber et al., 2016).

Down-regulation of miR-122 in liver tumours results in high expression of ABC transporter proteins and causes increased drug export of doxorubicin in liver cancer patients (Xu et al., 2011). Similarly, ABCB1 transporter expression is up-regulated in hepatocellular cancer cells when the post-transcriptional regulator miR-223 is down-regulated and the result is again resistance to doxorubicin treatment (Yang et al., 2013b).

Down-regulation of microRNAs let-7g and let-7i **results** in increased expression of ABCC10 that in turn is responsible for resistance to cisplatin therapy in oesophageal cancer patients (Wu et al., 2016a).

An important barrier for oral anticancer drugs is represented by intestinal epithelial cells of the GI tract (Ikemura et al., 2014; Peterson and Artis, 2014). The absorption of most nutrient components as well as drugs is related to a variety of influx transporters such as members of the SLC transporter family (Ikemura et al., 2014). The expression pattern of the SLC transporter varied according to the differentiation status of intestinal epithelial cells which is controlled by microRNAs (McKenna et al., 2010). Therefore, changes in the expression level of microRNAs have most probably an important influence on the drug up-take rate (McKenna et al., 2010). Up to now the role of microRNAs for the expression level of SLC transporter have been studied only in cell culture models for colon carcinoma, liver, pancreatic and gastric tumours (Dalmaso et al., 2011; Pullen et al., 2011). In colon cancer cells expression of miR-92b reduce the amount of SLC15A and SLC15A1 transporter resulting in decreased drug absorption (Dalmaso et al., 2011). In the context of liver and pancreatic tumours miR-29a, miR-29b and miR-124 target SLC16A1 and reduce the expression of this transporter (Pullen et al., 2011). Recently it was shown that miR-939 targets direct SLC34A2 in gastric cancer (Zhang et al., 2017). In 5-fluorouracil resistant gastric cancer miR-939 is down-regulated and results in increased expression level of SLC34A2. The transport protein SLC34A2 acts as mediator of miR-939 and activates the Ras/MEK/ERK pathway which is known to be deregulated often in cancer and to cause resistance to chemotherapy (Zhang et al., 2017). In *in-vitro* models of gastric cancer over-expression of miR-939 strongly decreased MEK1/2 phosphorylation as well as Raf-1 level, whereas SLC34A2 restoration rescued these effects (Zhang et al., 2017).

Also for some drug-metabolizing enzymes post-transcriptional regulations by miRNAs have been proven (Tsuchiya et al., 2006; Koturbash et al., 2012; Ikemura et al., 2014). **Due to their pivotal role in maintaining chemical and functional homeostasis of cells, cytochrome P450 enzymes are strictly controlled. Under physiological conditions, cytochrome P450 enzymes are involved in the regulation of endogenous molecules like bile acids and steroids and under pathological conditions in the case of chemotherapy these enzymes are important in regard to**

drug metabolism. De-regulated expression of cytochrome P450 enzymes is linked to drug resistance and therapy failure (Rendic and Guengerich, 2015).

For example, miR-378 targets mRNA coding for CYP2E1 and reduces the expression level of CYP2E1 protein in cell culture models of liver tumours (Mohri et al., 2010;Zhou et al., 2016). In liver cancer patients CYP2E1 expression is increased while miR-378 is down-regulated (Mohri et al., 2010;Zhou et al., 2016). Also, a direct regulation of CYP1B1 by miR-27b was demonstrated in hepatocellular cancer cell lines (An et al., 2017). Decreased expression of miR-27b results in high expression level of CYP1B1 and renders by this liver tumour resistant to docetaxel treatment (An et al., 2017).

In pancreatic cancer cells over-expression of miR-27b leads to down-regulation of CYP3A4 protein and results in drug resistance to cyclophosphamide because CYP3A4 is necessary for drug activation (Pan et al., 2009). MicroRNA-based regulation of enzymes involved in phase II reactions are less analysed but nevertheless, in the context of oesophageal cancer, regulation of glutathione S-transferase P1 (GSTP1) was found to be regulated by miR-133a (Kano et al., 2010). Reduced expression of the tumour suppressor miR-133a resulted in increased level of GSTP1 protein (Kano et al., 2010). In phase II detoxification reactions - including inactivation of platinum derivatives and alkylating reagents -GSTP1 catalyses the addition of glutathione to the drug activated during phase I reactions with electrophiles (Shea et al., 1988a;McLellan and Wolf, 1999a).

A more specific influence of non-coding RNAs on drug metabolism was demonstrated for 5-fluorouracil in liver and colon tumours (Offer et al., 2014;Chai et al., 2015). Dihydropyrimidine dehydrogenase, an important enzyme in 5-fluorouracil metabolism, is repressed by miR-494 in colon tumours and by miR-27a as well as miR-27b in liver cancer (Offer et al., 2014;Chai et al., 2015). The fact that the translation of one and the same enzyme in two different tissues is under the control of different miRNAs underlines the tissue-specific regulation and fine-tuning of protein expression that is exerted by miRNAs.

In liver cancer the translation of two of the most important targets of chemotherapeutic agents, dihydrofolate reductase and thymidylate synthase, are repressed by up-regulation of miR-215 (Wang et al., 2015b). Reduced expression of dihydrofolate reductase and thymidylate synthase leads to the development of insensitivity to doxorubicin treatment (Wang et al., 2015b).

Thymidylate synthase is the target of 5-fluoruracil therapy and this enzyme is down-regulated by increased expression of miR-192 and miR-215 in colon cancer patients (Boni et al., 2010). In this case altered microRNA expression results in down-modulation of the drug target and leads to therapy failure. In addition, miR-192 and miR-215 alter the cell-cycle control at multiple levels and prevent progression into the S-phase leading to 5-fluorouracil resistance (Boni et al., 2010).

A similar case was observed in pancreatic tumours where RRM2 (ribonucleotide reductase regulatory subunit M2) the target of gemcitabine is under direct control of miR-211 and let-7a (Bhutia et al., 2013;Maftouh et al., 2014). Decreased expression of miR-211 and let-7a results in higher RRM2 protein level and renders the tumours resistant to gemcitabine (Bhutia et al., 2013;Maftouh et al., 2014).

Deregulation of cell-cycle, DNA repair pathways and alteration in death pathways

Impaired cell cycle regulation and alteration of cell death pathways are common causes of drug resistance (Helleday et al., 2008;Rodrigues et al., 2012b). Increased cell cycle progression and reduced cell death rate lead to accumulation of mutations and uncontrolled cell proliferation, a hallmark of tumour cells (Hanahan and Weinberg, 2011). Errors in the DNA-damage response program pathways [nuclear excision repair (NER), base excision repair (BER), DNA mismatch repair (MMR)] play an important role in cancer progression and chemo-resistance (Hoeijmakers, 2001;Harper and Elledge, 2007;Jackson and Bartek, 2009;Pearl et al., 2015). A complex interaction interplay exists between non-coding RNAs and the DNA-damage pathways: on one hand the DNA-damage pathway induces the expression of several non-coding RNAs especially of microRNAs and on the other hand non-coding RNAs regulate directly the expression of several genes involved in DNA-damage pathway. This interaction is cell type specific and dependent on the intensity and nature of DNA damage (Pothof et al., 2009;Wouters et al., 2011;Chowdhury et al., 2013;Sharma and Misteli, 2013;Bottai et al., 2014).

LncRNA HOTAIR (HOX transcript antisense RNA) is highly expressed in a broad variety of solid tumours including liver, colorectal, pancreatic and gastrointestinal stromal tumours (Geng et al., 2011;Kogo et al., 2011;Niinuma et al., 2012). LncRNA HOTAIR reprogram

chromatin organization together with the polycomb repressive complex PRC2 (Kogo et al., 2011). Up-regulation of lncRNA HOTAIR results in higher expression level of members of the PRC2 complex (SUZ12, EZH2, and H3K27me3) (Kogo et al., 2011). Therefore, increased lncRNA HOTAIR expression is associated with a genome-wide reprogramming via PRC2 mediated epigenetic silencing of chromatin (Kogo et al., 2011). In addition lncRNA HOTAIR down-regulates cyclin-dependent kinase inhibitor 1 (p21(WAF/CIP1)) (Liu et al., 2013) causing the loss of an important regulator of the G₁ and S phase progression (el-Deiry et al., 1993;Waldman et al., 1995;Bunz et al., 1998). Due to the fact that p21(WAF/CIP1) represents a major target of p53 activity DNA damage in lncRNA HOTAIR expressing tumour cells don't go into cell cycle arrest and this promote cisplatin resistance (el-Deiry et al., 1993;Waldman et al., 1995;Bunz et al., 1998;Liu et al., 2013).

In oesophageal, gastric, colorectal and hepatocellular cancer as well as cholangiocarcinomas, lncRNA TUG1 (taurine-up-regulated gene 1) is involved in causing resistance to chemotherapy (Huang et al., 2015;Dong et al., 2016;Jiang et al., 2016;Li et al., 2016b;Wang et al., 2016a;Zhang et al., 2016a;Xu et al., 2017c). In tumour tissue lncRNA TUG1 is up-regulated and promotes cell growth by increased transcription of the *Bcl-2* gene and epigenetic silencing of cyclin-dependent protein kinase inhibitors (p15, p16, p21, p27 and p57) and pro-apoptotic genes (caspase-3, caspase-9 and Bax) (Huang et al., 2015;Dong et al., 2016;Jiang et al., 2016;Li et al., 2016b;Wang et al., 2016a;Zhang et al., 2016a;Xu et al., 2017c). Therefore, lncRNA TUG1 is an excellent example for the fact that non-coding RNAs target simultaneously the expression of different genes; beside increasing the expression level of the anti-apoptotic protein Bcl-2, expression of key players in the caspase-mediated apoptosis pathway are inhibited together with different cyclin-dependent protein kinase inhibitors. This results in decreasing the G₀/G₁ arrest during cell cycle and reduces the apoptosis rate of the tumour cells. Most probably lncRNA TUG1 has also a role in the epithelial-mesenchymal transition (Wang et al., 2016a;Xu et al., 2017c) that increases resistance to drug treatments further as outlined in detail below.

Also, the lncRNA PANDAR (promoter of CDKN1A antisense DNA damage-activated RNA) is often deregulated in different GI tumours like gastric, colorectal and hepatocellular cancer as well as cholangiocarcinoma (Peng and Fan, 2015;Ma et al., 2016a;Lu et al., 2017a;Xu et al., 2017b). In all these tumours up-regulation of lncRNA PANDAR results in increased

proliferation rate and reduced apoptosis (Peng and Fan, 2015;Ma et al., 2016a;Lu et al., 2017a;Xu et al., 2017b). LncRNA PANDAR interacts with the transcription factor NF- κ B, an important regulator for transcription of pro-apoptotic genes (Hung et al., 2011). This interaction between lncRNA PANDAR and NF- κ B results in decreased expression of pro-apoptotic genes and eventually leads to drug resistance (Peng and Fan, 2015;Ma et al., 2016a;Lu et al., 2017a;Xu et al., 2017b).

LncRNA UCA1 (urothelial carcinoma associated1) mediates resistance to doxorubicin treatment in gastric cancer (Shang et al., 2016). In *in-vitro* systems knockdown of lncRNA UCA1 overcomes the doxorubicin resistance due to an increased expression of PARP and reduced expression of Bcl-2 resulting in higher apoptosis rate (Shang et al., 2016).

Furthermore, it was shown that lncRNA UCA1 sequesters miR-204-5p in colorectal cancer and reduces the level of this microRNA in cancer cells (Bian et al., 2016). The consequence is enhanced cell proliferation and 5-fluorouracil resistance (Bian et al., 2016).

Another example of non-coding RNAs influencing cell-cycle is lncRNA ARA (adriamycin resistance associated) (Jiang et al., 2014;Cox and Weinman, 2016). LncRNA ARA was found to be over-expressed in doxorubicin resistant liver cancer cell lines compared to the parental cell lines (Jiang et al., 2014). Down-regulation of lncRNA ARA results in cell-cycle arrest in G2/M phase, suppressed proliferation, increased apoptotic cell death and, as expected, a reduced resistance against doxorubicin (Jiang et al., 2014;Cox and Weinman, 2016). Furthermore, lncRNA ARA is involved in the regulation of multiple signalling pathways including the MAPK-pathway (Jiang et al., 2014;Cox and Weinman, 2016). Beside lncRNA ARA the lncRNA URHC (up-regulated in hepatocellular carcinoma) is found among the most up-regulated lncRNAs in hepatocellular carcinoma. One target of lncRNA URHC is the tumour-suppressor ZAK (Xu et al., 2014b). Down-regulation of ZAK via lncRNA URHC results in increased cell proliferation and inhibits apoptosis (Xu et al., 2014b).

In pancreatic cancer lncRNA HOTTIP (HOXA transcript at the distal tip) up-regulates the homeobox-transcription factor HOX13 resulting in de-regulation of the cell cycle as well as gemcitabine resistance (Wang et al., 2011;Li et al., 2015e).

Down-regulation of lncRNA LOC285194 in oesophageal cancer results in resistance to chemoradiotherapy (radiation in combination with platinum- or paclitaxel-based

chemotherapy) by influencing cell-cycle progression and non-apoptotic cell death pathway via regulating VEGF receptor 1 (Tong et al., 2014).

In contrast, lncRNA MALAT-1 is strongly over-expressed in oesophageal tumour tissue and binds miR-107 and miR-217 (Lin and Xu, 2015; Wang et al., 2015c). MiR-107 and miR-217 decoy translates in reduced activity of the ATM-CHK2 signalling pathway leading to reduced cell-cycle arrest and cell death as response to DNA damage (Smith et al., 2010; Wang et al., 2015c) and over-expression of the transcription factor B-Myb – an important regulator for G1/S and G2/M cell-cycle progression and cell survival (Lin and Xu, 2015; Wang et al., 2015c).

In addition, several microRNAs have been identified as regulators for cell cycle progression and induction of cell death pathways. Therefore, deregulated microRNA expression pattern is often a reason for drug resistance in GI tumours.

Colorectal cancers with up-regulated mir-203 are resistant to oxaliplatin (Zhou et al., 2014). Failure of oxaliplatin therapy is caused by miR-203 mediated down-regulation of the important mediator protein for DNA damage response ATM (Zhou et al., 2014). As reaction to DNA damage, ATM induces the expression of DNA repair proteins, interrupts the cell cycle and induces cell death in the case of extended DNA damage (Choy and Watters, 2018). Oxaliplatin resistance can also be caused by up-regulation of miR-503-5p in colorectal cancer (Xu et al., 2017a). Increased expression of miR-503-5p results in down-regulation of the apoptotic protein PUMA (p53 upregulated modulator of apoptosis) and leads to resistance to oxaliplatin-induced apoptosis (Xu et al., 2017a). In colon cancer tissues down-regulation of miR-320 is linked to resistance to 5-fluorouracil therapy (Wan et al., 2015). Among the targets for miR-320 is the transcription factor SOX4 which is involved in inhibition of p53-mediated apoptosis as well as the cell cycle regulators FOXM1 and FOXQ1 both known to have oncogenic potential (Wan et al., 2015; Vishnubalaji et al., 2016).

In colorectal cancer cells miR-21 over-expression results in inhibition of the MMR proteins MSH2 and MSH6, two important proteins for DNA damage recognition and repair (Valeri et al., 2010a). Inhibition of MSH2 and MSH6 leads to reduced G2/M cell-cycle arrest caused by 5-fluorouracil induced DNA damage and lower apoptosis rate *in-vitro* and *in-vivo* (Valeri et al., 2010a). Therefore, miR-21 over-expression reduces the therapeutic efficacy of 5-fluorouracil-based chemotherapy in colorectal cancer treatment (Valeri et al., 2010a).

Furthermore, it was proven that the core mismatch repair proteins MSH2, MSH6 and MLH1 are also down-regulated by miR-155 potentially contributing to drug resistance (Valeri et al., 2010b). According to another study, 5-fluorouracil resistance in colorectal cancer cells can also be mediated by increased expression of miR-31 causing cell cycle deregulation and reduced apoptosis rate (Wang et al., 2010b; Cekaite et al., 2012). Efficacy of 5-fluorouracil treatment in colorectal cancer patients can also be limited due to up-regulation of anti-apoptotic proteins like XIAP (X-linked inhibitor of apoptosis) and UBE2N (ubiquitin-conjugating enzyme E2N) as a consequence of decreased miR-96 expression (Kim et al., 2015) or due to up-regulation of the anti-apoptotic proteins Bcl-2, Bcl-2-like protein 11 (BIM) or Bcl-2-like protein 2 (Bcl2L2) by reduced expression of miR-129, miR-10b or miR-195, respectively (Nishida et al., 2012; Karaayvaz et al., 2013; Qu et al., 2015). In other colon cancer studies reduced expression levels of miR-365, miR-1915 and miR-34a have been described as reason for increased expression of BCL-2 (Wang et al., 2010a; Nie et al., 2012; Xu et al., 2013).

Increased Bcl-2 expression has been identified as a reason for resistance to 5-fluorouracil in other GI tumours, too, but the posttranscriptional regulation of mRNA coding for Bcl-2 is under the control of different miRNAs; *e.g.* in gastric cancer diminished expression of miR-204 is the reason (Sacconi et al., 2012). According to another study up-regulation of Bcl-2 is caused by lower miR-15b and miR-16 expression level and leads to drug resistance in gastric cancer cells due to reduced apoptosis (Xia et al., 2008). MiR-25 over-expression was related to cisplatin resistance in gastric cancer cells (He et al., 2017). MiR-25 targets directly mRNAs coding for tumour suppressors like FOXO3a, ERBB2, FBXW7 (Zhao et al., 2014a; Gong et al., 2015; Li et al., 2015a; He et al., 2017). All these proteins are involved in cell cycle regulation and apoptosis (Huang and Tindall, 2006; Nho and Hergert, 2014; He et al., 2017). Up-regulation of miR-223 targets FBXW7 (F-box/WD repeat-containing protein 7) and leads to cell-cycle deregulation and cisplatin resistance in gastric tumours (Zhou et al., 2015). Furthermore, up-regulation of miR-103/107 results in decreased expression of caveolin-1 in gastric cancer cells (Zhang et al., 2015d). The tumour suppressor caveolin-1 is a counter regulator for the Ras-p42/p44 MAP kinase pathway and due to the down-regulation by miR-103/107 increased activity of the Ras-p42/44 Map kinase pathway results in increased cell cycle progression and reduced cell death (Le Gall et al., 2000; Mebratu and Tesfagzi, 2009). In gastric cancer increased cell cycle progression is also caused by

increased expression of miR-215 resulting in reduced expression of the tumour suppressor retinoblastoma 1, an important cell cycle regulator (Deng et al., 2014; Xu and Fan, 2015). Up-regulation of miR-106a targets FAS and inhibits the extrinsic apoptotic pathway in gastric cancer (Xiao et al., 2009; Wang et al., 2013c). In turn, reduced amount of FAS leads to increased cell proliferation, reduced apoptosis rate and drug resistance (Xiao et al., 2009; Wang et al., 2013c).

Over-expression of miR-21 inhibits cell cycle arrest resulting in increased cell proliferation, reduced apoptotic rate, gemcitabine and 5-fluorouracil resistance in pancreatic cancer (Moriyama et al., 2009; Park et al., 2009; Donahue et al., 2014). Similarly, in other pancreatic cancer studies, miR-21 over-expression results in reduced level of PTEN and Bcl-2 leading to activation of AKT-mTOR pathway, reduced apoptosis and resistance against gemcitabine treatment (Giovannetti et al., 2010; Dong et al., 2011). Increased expression of miR-214 represses directly ING4 in pancreatic tumour (Zhang et al., 2010). This impairs cell-cycle arrest, DNA repair as well as apoptosis and results in resistance to gemcitabine treatment (Zhang et al., 2010). The expression of the important pro-apoptotic protein BIM is reduced by miR-17-5p in pancreatic cancer and results in decreased apoptotic rate leading to resistance to gemcitabine treatment (Yan et al., 2012). Therapy failure is also caused by the repression of a tumour suppressor network involved in cell cycle and apoptosis regulation composed of PDCD4, BTG2 and NEDD4L by the combined action of miR-21, miR-23a and miR-27a (Frampton et al., 2014a; Frampton et al., 2014b). Furthermore, over-expression of miR-1246 results in decreased expression of cyclin-G2 and impairs the cell cycle regulation resulting in resistance to gemcitabine (Hasegawa et al., 2014). Recently miR-1307 was identified to be responsible for FOLFIRINOX resistance in pancreatic cancer (Carotenuto et al., 2018). MiR-1307 is up-regulated in *in-vitro* models of FOLFIRINOX resistant pancreatic cancer as well as in patient derived material compared to the surrounding tissue (Carotenuto et al., 2018). Reduced apoptosis rate and an extended acceptance of DNA damage seems to be the consequence of higher miR-1307 expression (Carotenuto et al., 2018).

In hepatocellular carcinoma the liver specific miR-122 is down-regulated and as consequence the expression of the target gene *CCNG1* is increased (Fornari et al., 2009). High level of cyclin G1 protein is found in several human tumours and results in reduced cell cycle control in the G2/M phase and modulation of p53 activity (Fornari et al., 2009; Xu et al., 2011). This

results in reduced DNA-repair and diminished apoptotic rate (Fornari et al., 2009; Xu et al., 2011). As already mentioned above, ABC transporter proteins are highly expressed in liver tumours due to the missing post-transcriptional regulator miR-122 (Xu et al., 2011). All these effects caused by miR-122 down-regulation promote doxorubicin resistance in liver cancer patients (Fornari et al., 2009; Xu et al., 2011). Another reason for doxorubicin resistance in liver cancer is based on reduced expression of miR-26b (Fan et al., 2008). Among the miR-26b targets in liver are the NF- κ B activating proteins TAB3 and TAK1 (Fan et al., 2008; Zhao et al., 2014b). Therefore, a reduced expression of miR-26b results in increased activation of NF- κ B and promotes drug resistance (Fan et al., 2008; Zhao et al., 2014b). Also, down-regulation of miR-101 is described as reason for resistance to doxorubicin in hepatocellular carcinoma (He et al., 2016). The anti-apoptotic protein Mcl-1 is among the targets of miR-101 and high levels of Mcl-1 renders liver tumour cells resistant to doxorubicin treatment (He et al., 2016). Furthermore, doxorubicin treatment failure in liver cancer patients has been connected to down-regulation of miR-199a-3p (Fornari et al., 2010). Besides targeting mTOR and c-Met, miR-199a-3p influences cell cycle regulation (Fornari et al., 2010). Decreased miR-199a-3p level results in down-regulation of the G1-checkpoint CDK inhibitors p21 (CDKN1A) and p27 (CDKN1B) and abrogate the G1 arrest following damage to DNA (Abukhdeir and Park, 2008; Fornari et al., 2010). In another study down-regulation of the G1 inhibitor CDKN1A in hepatocellular carcinoma was linked to up-regulation of miR-519d (Fornari et al., 2012). Consequently the apoptotic rate is reduced due to down-regulated miR-199a-3p as well as up-regulated miR-519d expression (Fornari et al., 2010; Fornari et al., 2012).

Another important tumour suppressor protein involved in resistance to anti-cancer drugs is PTEN because it is a main regulator for PI3K-AKT-mTOR pathway which is often hyperactivated in cancer and is one of the drivers for tumour growth and survival (Khan et al., 2013; LoRusso, 2016). PTEN itself is regulated by different microRNAs in different GI tumours, *e.g.* by miR-21 in liver and gastric cancer, miR-22 in p53-mutated colon cancer and mir-17-5p in colorectal cancer (Meng et al., 2007; Li et al., 2011; Zhang et al., 2012; Yang et al., 2013a; Fang et al., 2014). In all cases up-regulation of microRNAs results in decreased PTEN level in the tumour cell and subsequent activation of AKT-mTOR pathways resulting

in resistance to cisplatin (gastric cancer), paclitaxel (p53-mutated colon tumour) and FOLFOX (colorectal cancer) (Meng et al., 2007; Li et al., 2011; Zhang et al., 2012; Yang et al., 2013a; Fang et al., 2014). Down-regulation of PTEN due to over-expression of miR-19a and miR-19b in gastric cancer results in multi-drug resistance (Wang et al., 2013a).

Furthermore, mTOR is an important regulator under physiological as well as pathological conditions. In p53 mutant colorectal cancer mTOR is down-regulated by miR-338-3p and results in resistance to 5-fluorouracil treatment (Han et al., 2017). Indeed, inhibition of miR-338-3p in cell culture models restored sensitivity to 5-fluorouracil (Han et al., 2017) likely due to increased autophagy and reduced apoptosis following decrease in mTOR expression (Gonzalez et al., 2014; Han et al., 2017).

Autophagy is a further mechanism for chemoresistance (Song et al., 2009; Huang et al., 2016; Gozuacik et al., 2017; Xiong et al., 2017). In liver cancer up-regulation of lncRNA HULC activates autophagy by increasing the expression of ubiquitin-specific peptidase 22 (USP22) which in turn prevents the ubiquitin-mediated degradation of silent information regulator 1 (SIRT1) by removing the conjugated polyubiquitin chains from SIRT1 (Xiong et al., 2017). Autophagy causes resistance to oxaliplatin, 5-fluorouracil and epirubicin treatments in liver tumours (Xiong et al., 2017). In addition, lncRNA HULC down-regulates the expression of microRNAs that target directly the 3'-UTR of USP22 (miR-6825-5p, miR-6845-5p and miR-6886-3p) in liver cancer cells and prevents by this inhibition of USP22 at translational level (Xiong et al., 2017).

LncRNA MALAT-1 is highly expressed in gastric cancer cells resistant to 5-fluorouracil and cis-platin, respectively, compared to parental gastric cancer cells (YiRen et al., 2017). LncRNA MALAT-1 quenches miR-23b-3p and subsequently increases the expression of ATG12, an important regulator of autophagy (YiRen et al., 2017).

In oxaliplatin resistant colon cancer miR-409-3p is down-regulated so that the direct target Beclin-1 is expressed and induces autophagy (Tan et al., 2016). Over-expression of miR-409-3p results in low autophagic activity and overcomes oxaliplatin resistance in model systems of colon cancer (Tan et al., 2016).

Induction of epithelial-mesenchymal transition

Drug resistance can be caused by epithelial-mesenchymal transition (EMT) (Bedi et al., 2014;Heery et al., 2017). Several EMT-related signalling pathways are well known to be involved in mediating drug resistance in tumours (Nurwidya et al., 2012;Housman et al., 2014;Du and Shim, 2016;Heery et al., 2017). Cells undergoing EMT have several features in common with cancer stem cells (*e.g.* increased drug efflux pumps and anti-apoptotic effects) and furthermore EMT is instrumental for generation and maintenance of cancer stem cells (Housman et al., 2014;Du and Shim, 2016;Heery et al., 2017).

The lncRNA PVT1 (plasmacytoma variant translocation 1) has been found to be elevated in nearly all GI tumours including gastric, oesophageal, pancreatic, colon and liver cancers (Zheng et al., 2016;Wu et al., 2017;Zeng et al., 2017;Zhou et al., 2017). Increased expression of lncRNA PVT1 results in EMT and drug resistance (Zheng et al., 2016;Wu et al., 2017;Zhou et al., 2017).

The tumour suppressor lncRNA LEIGC prevents normal cells to undergo EMT. Therefore, the reduced expression of lncRNA LEIGC in gastric cancer fosters EMT and results in resistance to 5-fluorouracil treatment (Han et al., 2014b;Fang et al., 2015).

Up-regulation of lncRNA HULC has been correlated to induced EMT and suppressed apoptosis in gastric tumours leading to cisplatin resistance (Zhao et al., 2014c;Zhang et al., 2016b).

Increased expression of lncRNA-ATB (lncRNA-activated by TGF- β) in liver cancer results in competition with members of the miR-200 family for binding sites in the 3'-UTR of mRNAs coding for the transcription factors ZEB1 and ZEB2 (Yuan et al., 2014). In turn, high expression of ZEB1 and ZEB2 causes EMT and increased drug resistance (Yuan et al., 2014).

In pancreatic cancer the lncRNA MALAT-1 is a regulator of EMT (Ying et al., 2012;Jiao et al., 2014). In addition, the lncRNA MALAT-1 suppress G2/M cell cycle arrest and apoptosis leading to resistance to gemcitabine treatment (Jiao et al., 2014). As demonstrated by this example, the same lncRNA can induce resistance to chemotherapy by regulating different mechanisms at the same time.

Induction of EMT and resistance to gemcitabine treatment in pancreatic cancer cells can also be caused by miR-223 over-expression (Ma et al., 2015). Inhibition of miR-223 restored the

sensitivity of pancreatic cancer cell lines to gemcitabine treatment (Ma et al., 2015). Similarly, gemcitabine resistance in pancreatic cancer can also be caused by down-regulation of microRNAs as demonstrated for miR-200 (miR-200a, miR-200b and miR-200c) and let-7 family resulting in EMT (Li et al., 2009; Yu et al., 2010).

In colon cancer cells down-regulation of miR-147 results in EMT and increases the phosphorylation rate of AKT (Lee et al., 2014). Beside the activation of the PI3K-AKT pathway, the lower expression level of miR-147 also activates the TGF- β pathway and eventually leads to resistance to gefitinib treatment (Lee et al., 2014). Increased expression of miR-224 in colon cancer tissue was identified as another reason for resistance to 5-fluorouracil treatment. Increased miR-224 expression translates in increasing phosphorylation rate of extracellular signal-regulated kinase (ERK) and AKT, resulting in activation of both pathways (Amankwatia et al., 2015). In addition, miR-224 seems to activate also EGFR dependent- and NF- κ B-signalling pathway leading to EMT (Amankwatia et al., 2015).

Cancer cell stemness

A further reason for drug resistance is the presence of cancer stem cells. Cancer stem cells are well known for being refractory to chemotherapies and therefore cause therapy failure and tumour recurrence or progression (Reya et al., 2001; Ischenko et al., 2010; Li et al., 2010; Shankar et al., 2011; Srivastava et al., 2011; Nguyen et al., 2012; Pattabiraman and Weinberg, 2014). Once again non-coding RNAs especially lncRNAs and microRNAs are involved in sustaining the cancer stem cell niche (Tay et al., 2008; Liu and Tang, 2011; Sun et al., 2014; Garg, 2015; Chen et al., 2017).

The lncRNA UCA1 (urothelial carcinoma associated 1; identical with lncRNA CUDR (cancer up-regulated drug resistant)) is strongly expressed in different tumours; among these, gastric, hepatocellular, pancreatic, colorectal cancers and oesophageal squamous cell carcinoma (Han et al., 2014a; Li et al., 2014b; Wang et al., 2015a; Chen et al., 2016; Shang et al., 2016; Chen et al., 2017; Li et al., 2017; Wang et al., 2017). LncRNA UCA1 binds to several microRNAs in different tumours (e.g. miR-216b in liver cancer, miR-204 in oesophageal and colon cancer, miR-27b in gastric cancer) and influences entire transcriptional programs as well as response towards therapy (Wang et al., 2015a; Bian et al.,

2016;Fang et al., 2016b;Jiao et al., 2016;Wang et al., 2017). Well-established up-regulated targets of lncRNA UCA1 are members of the Wnt- β -catenin signalling pathway, several transcription factors and cell division regulators (Wang et al., 2008;Li and Chen, 2016). For cell stem cells the Wnt- β -catenin pathway is of pivotal importance for cell self-renewal and mediating drug resistance (Taipale and Beachy, 2001;Fan et al., 2014a). Over-expression of lncRNA UCA1 results in resistance to cancer treatments with tamoxifen, 5-fluorouracil, gemcitabine, cisplatin, doxorubicin, imatinib and tyrosine-kinase inhibitors targeting EGFR (Bian et al., 2016;Shang et al., 2016;Li et al., 2017;Wang et al., 2017).

Silencing of lncRNA UCA1 in *in-vitro* and *in-vivo* systems proved the oncogenic role of lncRNA UCA1 in gastric cancer (Shang et al., 2016;Li et al., 2017). Reduced expression level of lncRNA UCA1 results in reduced proliferation rate, increased apoptosis rate and overcomes the resistance to doxorubicin (Shang et al., 2016;Li et al., 2017). Furthermore, lncRNA UCA1 is a direct regulator of the PI3K-AKT-mTOR pathway (Li et al., 2017) which is often found to be deregulated in human cancers and is known to contribute to chemoresistance of cancer cells (Xia and Xu, 2015;Safa, 2016). In another study over-expression of lncRNA UCA1 was shown to cause reduced miR-27 expression causing diminished apoptosis of gastric cancer cells due to increased Bcl-2 protein level in combination with reduced cleaved caspase-3 (Fang et al., 2016b). This results in multidrug resistance of gastric tumours (Fang et al., 2016b).

Over-expression of lncRNA UCA1 is also a reason for chemo-resistance against 5-fluorouracil treatment in colon cancer (Bian et al., 2016). lncRNA UCA1 causes resistance by binding miR-204-5p and consequently up-regulating the expression of its target genes Bcl-2, RAB22A and CREB1 (Bian et al., 2016). MiR-21 was identified as an important player in regard to failure of 5-fluorouracil therapy in colon cancer patients (Yu et al., 2013). MiR-21 is able to increase the number of undifferentiated cancer stem cells during 5-fluorouracil treatment and contributes by this to therapy failure (Yu et al., 2013).

In liver cancer lncRNA UCA1 contributes to chemotherapy resistance and malignant transformation of hepatocyte-stem cells (Gui et al., 2015;Li and Chen, 2016;Li et al., 2016a;Chen et al., 2017;Huang et al., 2017;Zheng et al., 2017). lncRNA UCA1 increases directly the transcription rate of the oncogene c-myc well known to be involved in drug resistance as well as in activating stem-cell like properties in hepatocarcinoma (Walker et al.,

1996;Lin et al., 2007;Pyndiah et al., 2011;Akita et al., 2014;Pu et al., 2015). Furthermore, lncRNA UCA1 also induces the expression of lncRNA HULC (highly up-regulated in liver cancer) in liver cancer and lncRNA HULC in turn stimulates the activity of the Wnt- β -catenin pathway (Gui et al., 2015). In addition, lncRNA UCA1 forms a complex with the cell-cycle regulator cyclin-D which enhances the expression of lncRNA H19 by inhibiting the methylation of the lncRNA H19 promoter (Pu et al., 2015;Chen et al., 2017). High level of lncRNA H19 induces the telomerase activity and enhances the length of telomere thereby supporting the stem cell properties (Hiyama and Hiyama, 2007;Pu et al., 2015;Wu et al., 2016b). Another effect of lncRNA UCA1 is the enhanced phosphorylation of the tumour suppressor retinoblastoma protein 1(RB1). RB1 phosphorylation results in increased cell cycle progression and in interaction of the phosphorylated retinoblastoma protein 1 with the SET1A complex. Such interaction catalyses the transcription-activating methylation of histone H3 lysine-4 on several gene promoters including telomeric repeat-binding factor 2 promoter an important component for the telomerase extension process (Fang et al., 2016a;Li et al., 2016a).

In liver cancer as well as in pancreatic, gastric, oesophageal and colon cancers a critical role in inducing the transformation of stem cells into cancer stem cell has been demonstrated for lncRNA HOTAIR (Chen et al., 2013;Endo et al., 2013;Kim et al., 2013;He et al., 2014;Mohamadkhani, 2014;Li et al., 2015b;Chen et al., 2017). LncRNA HOTAIR is a strong activator for expression of *OCT4*, *RNF51*, *CD44* and *CD133* genes – all these proteins are involved in reprogramming the gene network to acquire cancer stem cell properties (Padua Alves et al., 2013;Zhu et al., 2014). LncRNA HOTAIR expression causes resistance against cisplatin and doxorubicin treatment in liver cancer model systems (Yang et al., 2011) and renders gastric tumours resistant to cisplatin therapy by binding miR-126 and activating the PI3K-AKT-mTOR pathway (Yan et al., 2016). In the context of several GI cancer stem cells it has been shown that lncRNA HOTAIR down-regulates the expression of histone methyltransferase SETD2 and reduces the phosphorylation rate of SETD2 resulting in reduced trimethylation of histone H3 lysine-36 on several gene promoter, e.g. Wnt inhibitory factor-1 (WIF-1) (Ge et al., 2013;Kim et al., 2013;Ding et al., 2014;Li et al., 2015b). Reduced WIF-1 expression leads to activation and increased signalling through the Wnt- β -catenin pathway (Ge et al., 2013;Kim et al., 2013). Furthermore, the modulated chromatin organisation account for a reduced efficiency of the mismatch repair system and damaged

DNA can escape from corrections leading to microsatellite instability (MSI) and altered expression of cell cycle regulators as well as reduced apoptosis (Gupta et al., 2010;Valeri et al., 2010b;Chen et al., 2013;Li et al., 2013;Li et al., 2015b). In addition, lncRNA HOTAIR induces accumulation of replication errors by hindering the complex formation of MSH2 with MSH6; one essential dimer for DNA mismatch recognition and repair (Yang et al., 2004;Valeri et al., 2010a;Valeri et al., 2010b;Edelbrock et al., 2013;Pfister et al., 2014).

In pancreatic cancer the oncogenic lncRNA MALAT-1 (metastasis-associated lung adenocarcinoma transcript-1) contributes to the expression of the cancer stem cell marker CD133, CD44, CD24 and aldehyde-dehydrogenase (Fan et al., 2014b;Jiao et al., 2014;Jiao et al., 2015). In addition, the expression of the core pluripotent factors OCT4, NANOG and SOX2 are also under the control of lncRNA MALAT-1 (Jiao et al., 2015). LncRNA linc-ROR inhibits the expression of p53 and activates by this the transcription factor ZEB1 in pancreatic cancer (Wellner et al., 2009). ZEB1 in turn suppress the expression of the miR-200 family that leads to maintenance of pancreatic cancer stemness and induces EMT known to be responsible for paclitaxel resistance in pancreatic cancer patients (Wellner et al., 2009;Kim, 2017). Down-regulation of miR-205 results in increased expression of stem cell markers OKT3, OKT8 and CD44 in pancreatic cancer tissue and is linked to gemcitabine resistance (Singh et al., 2013). Re-expression of miR-205 is able to overcome the gemcitabine resistance in pancreatic cancer model systems (Singh et al., 2013).

The lncRNA-34a mediates an increase in self-renewal of colon cancer stem cells and induce Wnt as well as NOTCH signalling pathways via sequester miR-34a expression (Bu et al., 2013;Evans et al., 2015).

In hepatocellular carcinoma the lncRNA linc-ROR (long intergenic ncRNA regulator of reprogramming) is involved in regulating core pluripotent factors (OCT-4, NANOG, SOX2) necessary for the stem cell like phenotype and causes resistance to chemotherapy (Takahashi et al., 2014). LncRNA linc-ROR competes with miR-145 for the same binding sites present in the mRNAs coding for OCT-4, NANOG and SOX2 (Wang et al., 2013b). Presence of lncRNA linc-ROR prevents the binding of miR-145 to the mRNA of the core pluripotent factors resulting in translation of these mRNAs and maintains the stem cell phenotype (Wang et al., 2013b). Furthermore, the expression of CD133, another cancer stem cell marker, is directly induced by lncRNA linc-ROR (Takahashi et al., 2014).

MiR-130b is connected to cancer stem cells growth in liver tumours (Ma et al., 2010). Increased expression of miR-130b targets directly the mRNA coding for tumour protein 53-induced nuclear protein 1 and reduces the expression level of the corresponding protein (Ma et al., 2010). Furthermore, high level of miR-130b renders liver tumour cells resistant to doxorubicin treatment (Ma et al., 2010). Another reason for doxorubicin resistance in liver cancer patients is down-regulation of the tumour suppressor miR-101 resulting in increased protein expression of enhancer of zeste homolog 2 (EZH2) (Sasaki et al., 2008; Xu et al., 2014a). EZH2 is a histone-lysine N-methyltransferase enzyme that silence Wnt-pathway antagonists and other tumour suppressor genes on the transcriptional level by histone methylation (Cheng et al., 2011). Over-expression of EZH2 is positively correlated with increased Wnt- β -catenin signalling (Cheng et al., 2011).

MiR-221 is over-expressed in 5-fluorouracil resistant oesophageal tumours (Wang et al., 2016b). The mechanisms of resistance is mediated via down-regulation of the direct target DDK2 (dickkopf-related protein 2) and subsequent activation of the Wnt- β -catenin pathway (Wang et al., 2016b). Furthermore, increased miR-221 expression fosters EMT and facilitates the formation of tumour stem cells (Wang et al., 2016b).

In colon cancer stem cells, miR-451 was found to be down-regulated compared to colon cancer cells (Bitarte et al., 2011). Reduced level of miR-451 seems to be essential for the self-renewal of colon cancer stem cells (Bitarte et al., 2011). In addition, expression of ABCB1 transporter is increased in colon cancer stem cells due to lack of miR-451 post-transcriptional down-regulation resulting in resistance to irinotecan treatment (Bitarte et al., 2011).

MiR-1182 is often down-regulated in gastric cancer tissue (Zhang et al., 2015a). One direct target of miR-1182 is telomerase reverse transcriptase (hTERT), an enzyme that is involved in controlling the length of telomere. Over-expression of hTERT due to missing transcriptional regulation by miR-1182, results in cell immortality and stem-cell property of gastric cancer cells (Zhang et al., 2015a).

Targeted therapies and drug resistance

For GI cancer several targeted therapies exist (Table 3) (Jonker et al., 2007;Weber and McCormack, 2008;Loupakis et al., 2010;Roukos, 2010;Grothey et al., 2013;Muro et al., 2015;King et al., 2017). They are used alone or in combination with chemotherapy. Unfortunately in most cases the patients develop resistance also against these targeted therapies and the above outlined general principles of drug resistance based on non-coding RNA dysregulation are involved. Beside that non-coding RNAs interfering with the targeted protein itself or (up-)regulating the targeted signal pathway are involved in drug resistance (Roukos, 2010). Furthermore, therapy failure can be related to activation of alternative signal pathways by non-coding RNAs (Roukos, 2010;Lu et al., 2017b).

Recently it was demonstrated that resistance to cetuximab in colon cancer patients and in *in-vitro* 3-D-cell culture models can be caused by over-expression of lncRNA MIR100HG (Lu et al., 2017b). Two microRNAs, miR-100 and miR-125b, are generated from lncRNA MIR100HG and these microRNAs down-regulate in a concerted way five negative regulators of the Wnt/ β -catenin pathway resulting in increased Wnt signalling (Lu et al., 2017b). This kind of cetuximab resistance can be overcome by inhibition of Wnt signalling, underscoring the potential clinical relevance of the interactions between EGFR and Wnt/ β -catenin pathways (Lu et al., 2017b). Increased mir-125b expression is also correlated with trastuzumab resistance in HER2-positive gastric cancer patients but up to now the molecular basis for this resistance is unclear (Sui et al., 2017). Sorafenib resistance in hepatocellular carcinoma is caused by lncRNA TUC338 (Jin et al., 2017). RASAL-1 (RAS protein activator like-1) is a direct target of lncRNA TUC338 and high expression of lncRNA TUC338 inhibits the RASAL-1 expression resulting in activation of RAS-signalling (Jin et al., 2017). According to another *in-vitro* study, reduced expression of miR-193b leads to higher expression of the anti-apoptotic protein Mcl-1 and renders hepatocellular carcinoma cells resistant to sorafenib treatment (Braconi et al., 2010).

Non-coding RNAs as potential biomarkers of resistance and novel therapeutics: promises and hurdles

Our review summarises most of the current evidence supporting the role of non-coding RNAs in resistance to chemotherapy and targeted agents. It is likely that, in the near future, given the promising and exciting results obtained with the use of immunotherapy in

gastroesophageal (Kang et al., 2017) and colorectal cancer (Le et al., 2017;Overman et al., 2018), new data will emerge on the already known regulation of PD-1, PD-L1 and CTLA-4 by non-coding RNAs and response to nivolumab and pembrolizumab (Cortez et al., 2016;Xu et al., 2016;Smolle et al., 2017).

The contribution of non-coding RNAs in resistance mechanisms to a broad range of anti-cancer treatments makes their use as biomarkers or novel therapeutics quite promising but several challenges remain.

Given microRNAs and, to a lesser extent, other non-coding RNAs can be reliably detected in tissues and bio-fluids such as plasma, serum and urine, it is tempting to hypothesize the use of non-coding RNA based tools to predict and monitor resistance to anticancer treatments. Few studies have already tested the validity of microRNAs as biomarkers of response to anticancer treatment in other cancers such as prostate (Lin et al., 2017), chronic lymphocytic leukaemia (Gagez et al., 2017) and sarcomas (Wiemer et al., 2017). In colorectal cancer, we (Sclafani et al., 2015) and others (Graziano et al., 2010;Zhang et al., 2011;Sha et al., 2014) have tested the contribution of a single nucleotide polymorphism (SNP) in the binding site of let-7 in the *KRAS* 3'UTR in predicting benefit from anti-EGFR treatment with conflicting results across different trials. Despite the good reproducibility of the assay, the predictive value of the test was not confirmed in all trials likely due to use of cetuximab in different context (neo-adjuvant, adjuvant and metastatic colorectal cancer, respectively). Similarly the analysis of a SNP in miR-608 led to contradicting results in patients treated with neo-adjuvant or adjuvant chemo- and radiochemo-therapy in colon and rectal cancers highlighting some of the challenges in validating data obtained in retrospective series (Lin et al., 2012;Xing et al., 2012;Pardini et al., 2015;Sclafani et al., 2016). Tissue (cancer *versus* stroma) and organ (colon *versus* rectum) specificity in non-coding RNA expression might represent potential explanations for different findings obtained in some of these studies. Beside SNPs, expression of microRNAs can be detected in fresh frozen or formalin fixed paraffin embedded tissues and serve as potential biomarker of sensitivity or resistance to treatment. Robust data have emerged from the retrospective analysis of a prospective phase III clinical trial (Laurent-Puig et al., 2016). In this study, *KRAS* wild-type patients were classified based on high or low miR-31-3p expression: patients with high expression were resistant to cetuximab while patient with low expression had good and durable responses

which translated in survival benefit. The miR-31 expression cut-off for the classification into high or low expression was predefined in the above study. However, one of the key challenges in validating these interesting findings will be design of a clinically approved assay that can accurately assign patients into one of these two categories. In this prospective, the use of different sources of material (i.e. primary colorectal cancer *versus* metastasis) might result in different basal expression of the microRNA and as such different scoring. Source of material and choice of reference controls represent important obstacles that might bias the definition of a threshold for high or low expression of microRNAs in tissues and bio-fluids. MicroRNAs can be detected in plasma, serum and urine samples and have been used for early detection and prognostic purposes in gastrointestinal cancer (Schultz et al., 2014; Shigeyasu et al., 2017; Ozawa et al., 2018). The use of digital droplet approaches allows the quantitative detection of copies of the microRNA of interest based on the starting volume of bio-fluids and, potentially overcomes or at least mitigates, the issues related to the normalization of data against reference controls, making the definition of cut-off easier to standardize. One study has reported the potential role of miR-126 in predicting and tracking response to chemotherapy and anti-VEGF treatment in colorectal cancer (Hansen et al., 2015) and, with the advent of digital quantitative technologies, more studies are expected.

In consideration of their role in cancer initiation, progression and resistance to treatment, non-coding RNAs and among them microRNAs have been proposed as potential therapeutics (Adams et al., 2017). A large body of pre-clinical evidence is available on the use of anti-microRNAs or molecules re-expressing microRNAs alone or in combination with other agents in order to increase efficacy and prevent or revert drug resistance (Rupaimoole and Slack, 2017). Inhibition of microRNAs has been tested in clinical trials in the context of HCV infection (Janssen et al., 2013; van der Ree et al., 2017) and in mesothelioma (van Zandwijk et al., 2017). These trials highlighted a huge potential for microRNA-based therapeutics but at the same time pinpointed some of the criticalities in further clinical development of such approaches. MiR-122 inhibition led to durable viral load reduction in both HCV trials and was associated with manageable side effects. Similarly, in mesothelioma patients treated with miR-16-loaded minicells the disease control rate was satisfactory and the toxicity profile acceptable warranting further investigations. Overall in both approaches the risk of off-target effects represent the main hurdle to be taken into account: indeed miR-122 inhibition has been associated with risk of developing liver cancer in pre-clinical models

(Hsu et al., 2012) and, similarly, over-expression of miR-16 might lead to uncontrolled cardiac effects as proven in the phase I trial (van Zandwijk et al., 2017). These effects might be increased in combination studies in which anti-microRNAs or microRNA-conjugates are delivered together with chemotherapy leading to cumulative side effects. Therefore a robust understanding of the biology underpinning microRNA deregulation in physiology and pathological conditions in order to implement effort that can minimise the risk of serious adverse events hampering the clinical development of microRNA-based strategies.

Conclusion:

Non-coding RNAs especially lncRNAs and microRNAs are important mediators for drug resistance. They function in an organ and tissue specific manner and through different molecular mechanisms. One non-coding RNA always have several targets and in the end deregulation of one non-coding RNA alters the expression level of several proteins in a tissue specific way. For example, in the case of miR-374b more than 700 genes have been identified as direct target in pancreatic tissue (Schreiber et al., 2016). Drug resistance is a dynamic process caused by several cell and non-cell autonomous mechanisms. Given non-coding RNAs can simultaneously control several cancer-associated pathways, non-coding RNA dysregulation plays a crucial role in treatment resistance. Future studies will continue to shed insights in the fine interplay among lncRNA, microRNA and their target genes and might provide opportunities for more effective strategies to prevent or overcome resistance. In the interim, given non-coding RNAs and especially microRNAs can be tested in tissues and biofluids in a rapid, cost/effective and robust way. More investigational studies should explore their utility to monitor and forecast treatment response and resistance in order to personalise treatments and improve patient's outcomes.

Conflict of Interest:

No conflicts to declare.

Author Contributions:

853 NV and JCH: idea, conception and writing the review.

854

In review

Figure legends:

Figure 1: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in liver cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 2: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in oesophageal cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 3: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in gastric cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 4: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in colon and colorectal cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 5: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in pancreatic cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 6: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in gastrointestinal stromal cancer. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Figure 7: Role of non-coding RNAs for the different reasons that can cause resistance to anticancer drugs in cholangiocarcinoma. For details about target genes and regulated protein expression by the non-coding RNAs see text.

Table1: Overview about non-coding RNAs involved in resistance to anticancer drugs in gastrointestinal tumours.

Abbreviation used: GI=gastrointestinal; vRNA=vault RNA; lncRNA=long non-coding RNA; miR=microRNA; EMT=epithelial-mesenchymal transition

Non-coding RNA	GI cancer type	Causing drug resistance via	Reference
lncRNA AK022798	gastric cancer	increasing the expression of <i>ABCB1</i> gene	(Hang et al., 2015)
lncRNA ANRIL	gastric cancer	increasing the expression of <i>MDR1</i> gene	(Zhang et al., 2015c; Lan et al., 2016)
lncRNA ARA	liver cancer	reduced G2/M cell-cycle arrest; reduced apoptosis rate; de-regulation of MAPK-pathway	(Jiang et al., 2014; Cox and Weinman, 2016)
lncRNA-ATB	liver cancer	increased expression of ZEB1 and ZEB2; induced EMT	(Yuan et al., 2014)
lncRNA CCAL	colorectal cancer	increasing the expression of <i>ABCB1</i> gene; increased activity of Wnt/ β -catenin pathway	(Ma et al., 2016b)
lncRNA H19	liver cancer oesophageal cancer	up-regulation of membrane glycoprotein p95; elevating the expression of <i>MDR1</i> gene by increasing promoter methylation; increasing telomere length	(Hiyama and Hiyama, 2007; Tsang and Kwok, 2007; Matouk et al., 2013)
lncRNA HOTAIR	liver cancer colorectal cancer pancreatic cancer GI stromal tumour	increased expression of PRC2 complex members; genome-wide changes in transcription process due to epigenetic chromatin silencing; down-regulation of p21(WAF/CIP1); repression of G1/S cell-cycle arrest; increased proliferation rate; reduced DNA-damage response	(el-Deiry et al., 1993; Geng et al., 2011; Kogo et al., 2011; Liu et al., 2013)
lncRNA HOTAIR	colon cancer pancreatic cancer gastric cancer	transformation of stem cells into cancer stem cells due to activation of <i>OCT4</i> , <i>RNF51</i> ,	(Yang et al., 2004; Edelbrock et al., 2013; Ge et al.,

Non-coding RNAs and anticancer drugs

	oesophageal cancer	<i>CD44</i> and <i>CD133</i> gene expression; increased activity of Wnt/ β -catenin pathway; modulation of chromatin organisation leads to reduced efficiency of the mismatch repair system; increased MSI; reduced apoptosis rate; inhibition of the expression of miR-126 and activating the PI3K-AKT-mTOR pathway (in gastric cancer)	2013;Kim et al., 2013;Padua Alves et al., 2013;Zhu et al., 2014;Yan et al., 2016)
lncRNA HOTTIP	pancreatic cancer	increased expression of transcription factor HOX13; cell cycle deregulation	(Wang et al., 2011;Li et al., 2015e)
lncRNA HULC	liver cancer	increased activity of Wnt- β -catenin; increased expression of USP22 and SIRT1; reduced expression of miR-6825-5p, miR-6845-5p, miR-6886-3p; increased autophagy pathway	(Xiong et al., 2017)
lncRNA HULC	gastric cancer	induced EMT; suppressed apoptosis	(Zhao et al., 2014c;Zhang et al., 2016b)
lncRNA LEIGG	gastric cancer	induced EMT	(Han et al., 2014b;Fang et al., 2015)
lncRNA linc-ROR	pancreatic cancer	inhibition of p53; inhibition of the expression of miR-200 family; increased expression of the transcription factor ZEB1; induced EMT	(Wellner et al., 2009;Kim, 2017)
lncRNA linc-ROR	liver cancer	preventing the binding of miR-145 to pluripotent factors OKT-4, NANOG and SOX2 resulting in increased expression of these transcription factors necessary for sustain stem cell character	(Wang et al., 2013b;Takahashi et al., 2014)
lncRNA LOC285194	oesophageal cancer	cell-cycle deregulation; blocking non-apoptotic cell death pathway	(Tong et al., 2014)

Non-coding RNAs and anticancer drugs

lncRNA MALAT-1	oesophageal tumour	binds miR-107 and miR-217; reduced activity of the ATM-CHK2 signalling pathway; reduced cell-cycle arrest and cell death as response to DNA damage; increased expression of transcription factor B-Myb	(Smith et al., 2010; Lin and Xu, 2015; Wang et al., 2015c)
lncRNA MALAT-1	pancreatic cancer	increased expression of cancer stem cell marker CD133; increased expression of pluripotent factors OCT4, NANOG and SOX2; induced EMT; repression of G2/M cell-cycle arrest; reduced apoptosis rate	(Ying et al., 2012; Jiao et al., 2014; Jiao et al., 2015)
lncRNA MALAT-1	gastric cancer	sequestering of miR-23b-3p; increased expression of ATG12; increased autophagy	(YiRen et al., 2017)
lncRNA MIR100HG	colon cancer	increased activity of Wnt- β -catenin pathway	(Lu et al., 2017b)
lncRNA MRUL	gastric cancer	increasing the expression of <i>MDR1</i> gene	(Wang et al., 2014)
lncRNA PANDAR	gastric cancer colorectal cancer hepatocellular cancer cholangiocarcinoma	interacts with the transcription factor NF- κ A resulting in reduced translation of pro-apoptotic genes – leading to reduced apoptosis rate and increased proliferation	(Hung et al., 2011; Peng and Fan, 2015; Ma et al., 2016a; Lu et al., 2017a; Xu et al., 2017b)
lncRNA PVT1	gastric cancer oesophageal cancer pancreatic cancer colon cancer liver cancer	induced EMT	(Zheng et al., 2016; Wu et al., 2017; Zhou et al., 2017)
lncRNA PVT-1	gastric cancer	increasing the expression of <i>MDR1</i> gene	(Zhang et al., 2015c; Lan et al., 2016)
lncRNA TUC338	hepatocellular cancer	inhibiting the RASAL-1 pathway	(Jin et al., 2017)
lncRNA TUG1	oesophageal cancer gastric cancer colorectal cancer hepatocellular cancer cholangiocarcinoma	increasing the expression of <i>Bcl-2</i> gene; reducing the expression of cyclin-dependent protein kinase, caspase-3, caspase-9 and Bax; decreasing G0/G1 arrest during cell cycle; reducing apoptosis rate; inducing EMT	(Huang et al., 2015; Dong et al., 2016; Jiang et al., 2016; Li et al., 2016b; Wang et al., 2016a; Zhang et al., 2016a; Xu et al., 2017c)

Non-coding RNAs and anticancer drugs

lncRNA UCA1 (identical with lncRNA CDUR)	liver cancer colorectal cancer pancreatic cancer gastric cancer oesophageal cancer	sequestering microRNAs (miR-216b in liver cancer; miR-204-5p in colorectal and oesophageal cancer; miR-27 in gastric cancer); increase expression of lncRNAs (HULC; H19); increased activity of Wnt- β -catenin pathway; increased activity of PI3K-AKT-mTOR pathway; increased phosphorylation of tumour suppressor retinoblastoma; increased expression of c-myc; increased cell-cycle progression; increased expression of anti-apoptotic protein Bcl-2; reduced expression of PARP (in gastric cancer); reduced apoptosis rate. In liver cancer additional effects: transformation of stem cells into cancer stem cells due to increased c-myc expression; increasing telomere length	(Walker et al., 1996; Hiyama and Hiyama, 2007; Wang et al., 2008; Gui et al., 2015; Pu et al., 2015; Bian et al., 2016; Fang et al., 2016a; Fang et al., 2016b; Li and Chen, 2016; Shang et al., 2016; Chen et al., 2017; Li et al., 2017)
lncRNA URHC	liver cancer	reduced expression of the tumour suppressor ZAK; increased proliferation rate; reduced apoptosis rate	(Xu et al., 2014b)
lncRNA-34a	colon cancer	increased activity of Wnt- β - catenin pathway; increased activity of NOTCH pathway; increasing the self-renewal of cancer stem cells	(Bu et al., 2013; Evans et al., 2015)
miR let-7 family	pancreatic cancer	induced EMT	(Li et al., 2009)
miR let-7a	pancreatic tumours	increased expression of RRM2	(Bhutia et al., 2013)
miR let-7g	oesophageal cancer	increased expression of ABCC10	(Wu et al., 2016a)
miR let-7i	oesophageal cancer	increased expression of ABCC10	(Wu et al., 2016a)
miR-100	colon cancer	increased activity of Wnt- β - catenin pathway	(Lu et al., 2017b)
miR-101	liver cancer	increased expression of EZH2; increased activity of Wnt- β -catenin pathway; increased expression of Mcl-	(Sasaki et al., 2008; Xu et al., 2014a; He et al., 2016)

Non-coding RNAs and anticancer drugs

		1; reduced apoptosis rate	
miR-10b	colorectal cancer	increased expression of anti-apoptotic protein BIm	(Nishida et al., 2012)
miR-103/107	gastric cancer	reduced expression of tumour-suppressor caveolin-1; activation of Ras-p42/p44 MAP pathway; reduced apoptosis rate	(Le Gall et al., 2000; Mebratu and Tesfagzi, 2009; Zhang et al., 2015d)
miR-106a	gastric cancer	reduced expression of FAS; reduced apoptosis rate	(Xiao et al., 2009; Wang et al., 2013c)
miR-1182	gastric cancer	increased expression of hTERT	(Zhang et al., 2015a)
miR-122	liver cancer	increased expression of ABC proteins; increased expression of cyclin G1; reduced G2/M cell-cycle arrest; reduced DNA repair; reduced apoptosis rate	(Fornari et al., 2009; Xu et al., 2011)
miR-124	pancreatic cancer liver cancer	reduced expression of SLC16A1	(Pullen et al., 2011)
miR-125b	colon cancer	increased activity of Wnt- β -catenin pathway	(Lu et al., 2017b)
miR-1246	pancreatic cancer	reduced expression of cyclin-G2; de-regulated cell-cycle	(Hasegawa et al., 2014)
miR-129	colorectal cancer	increased expression of anti-apoptotic protein Bcl-2	(Karaayvaz et al., 2013)
miR-1291	pancreatic cancer	increased expression of ABCC1	(Pan et al., 2013)
miR-130b	liver cancer	reduce expression of tumour protein 53-induced nuclear protein 1	(Ma et al., 2010)
miR-1307	pancreatic cancer	reduced apoptosis rate	(Carotenuto et al., 2018)
miR-133a	oesophageal cancer	increased expression of GSTP1	(Kano et al., 2010)
miR-145	colon carcinoma	increased expression of ABCB1	(Ikemura et al., 2013)
miR-147	colon cancer	induced EMT; increased phosphorylation of AKT; increased activity of PI3K-AKT-mTOR pathway; increased activity of TGF- β pathway	(Lee et al., 2014)

Non-coding RNAs and anticancer drugs

miR-155	colorectal cancer	inhibition of MSH2, MSH6 and MLH1	(Valeri et al., 2010b)
miR-15b	gastric cancer	increased expression of anti-apoptotic protein Bcl-2	(Xia et al., 2008)
miR-16	gastric cancer	increased expression of anti-apoptotic protein Bcl-2	(Xia et al., 2008)
mir-17-5p	colorectal cancer	reduced expression of PTEN expression; activation of AKT-mTOR pathways	(Fang et al., 2014)
miR-17-5p	pancreatic cancer	reduced expression of BIM	(Yan et al., 2012)
miR-1915	colon cancer	increased expression of BCL-2	(Xu et al., 2013)
miR-192	colon cancer	reduced expression of thymidylate synthase; altered cell-cycle control at multiple levels; prevent progression into the S-phase	(Boni et al., 2010)
miR-193b	hepatocellular cancer	increased expression of Mcl-1	(Braconi et al., 2010)
miR-195	colorectal cancer	increased expression of anti-apoptotic protein Bcl-2L2	(Qu et al., 2015)
miR-199a-3p	liver cancer	reduced G1/S cell-cycle arrest; increased expression of mTOR and c-Met; reduced apoptosis rate	(Abukhdeir and Park, 2008; Fornari et al., 2010)
miR-19a	gastric cancer	reduced expression of PTEN expression; activation of AKT-mTOR pathways	(Wang et al., 2013a)
miR-19b	gastric cancer	reduced expression of PTEN expression; activation of AKT-mTOR pathways	(Wang et al., 2013a)
miR-200a	pancreatic cancer	induced EMT	(Li et al., 2009)
miR-200b	pancreatic cancer	induced EMT	(Li et al., 2009)
miR-200c	pancreatic cancer	induced EMT	(Li et al., 2009; Yu et al., 2010)
miR-203	colorectal cancer	reduced expression of ATM; impaired DNA repair; reduced apoptosis rate	(Zhou et al., 2014)
miR-205	pancreatic cancer	increased expression of pluripotent factors OKT3, OKT8 and CD44	(Singh et al., 2013)
miR-21	colorectal cancer	inhibition of MSH2 and MSH6; reduced G2/M cell-cycle arrest; reduced apoptosis rate; increasing the number of undifferentiated cancer stem cells	(Valeri et al., 2010a; Yu et al., 2013)

Non-coding RNAs and anticancer drugs

miR-21	pancreatic cancer	reduced cell-cycle arrest; reduced expression of PTEN; activation of AKT-mTOR pathway; increased expression of anti-apoptotic protein Bcl-2; increased cell proliferation; reduced apoptosis rate	(Giovannetti et al., 2010; Dong et al., 2011)
miR-21	liver cancer gastric cancer	reduced expression of PTEN expression; activation of AKT-mTOR pathways	(Meng et al., 2007; Zhang et al., 2012; Yang et al., 2013a)
synergistic action of miR-21 miR-23a miR-27a	pancreatic cancer	reduced expression of the tumour suppressors PDCD4, BTG2 and NEDD4L; de-regulated cell-cycle; reduced apoptosis rate	(Frampton et al., 2014a; Frampton et al., 2014b)
miR-211	pancreatic tumours	increased expression of RRM2	(Maftouh et al., 2014)
miR-215	liver cancer	reduced expression of dihydrofolate reductase; reduced expression of thymidylate synthase	(Wang et al., 2015b)
miR-215	colon cancer	reduced expression of thymidylate synthase; altered cell-cycle control at multiple levels; prevent progression into the S-phase	(Boni et al., 2010)
miR-215	gastric cancer	reduced expression of retinoblastoma 1; altered cell-cycle control	(Deng et al., 2014; Xu and Fan, 2015)
miR-22	p53-mutated colon cancer	reduced expression of PTEN expression; activation of AKT-mTOR pathways	
miR-221	oesophageal cancer	reduced expression of DDK2; activation of Wnt/ β -catenin pathway; induced EMT	(Li et al., 2011; Wang et al., 2016b)
miR-223	liver cancer	increased expression of ABCB1	
miR-223	pancreatic cancer	induced EMT	(Ma et al., 2015)
miR-223	gastric cancer	reduced expression of FBXW7; altered cell-cycle control	(Zhou et al., 2015)
miR-224	colon cancer	induced EMT; increased phosphorylation of AKT und ERK; increased activity of PI3K-AKT-mTOR pathway; increased activity of ERK	(Amankwatia et al., 2015)

Non-coding RNAs and anticancer drugs

		pathway; activation of NF- κ B and EGFR dependent pathways	
miR-23a	microsatellite instable colon cancer	increased expression of ABCF1	(Li et al., 2015c)
miR-25	gastric cancer	reduced expression of FOXO3a, ERBB2 and FBXW7; cell-cycle deregulation; reduced apoptosis rate	(Zhao et al., 2014a; Gong et al., 2015; Li et al., 2015a; He et al., 2017)
miR-26b	liver cancer	increased activation of NF- κ B	(Fan et al., 2008; Zhao et al., 2014b)
miR-27a	liver cancer	reduced expression of dihydropyrimidine dehydrogenase	(Offer et al., 2014)
miR-27b	liver cancer	increased expression of CYP1B1; reduced expression of dihydropyrimidine dehydrogenase	(Offer et al., 2014; An et al., 2017)
miR-27b	pancreatic cancer	reduced expression of CYP3A4– resulting in cyclophosphamide resistance due to missing drug activation	(Pan et al., 2009)
miR-297	colorectal cancer	increased expression of ABCC2	(Xu et al., 2012)
miR-29a	pancreatic cancer liver cancer	reduced expression of SLC16A1	(Pullen et al., 2011)
miR-29b	pancreatic cancer liver cancer	reduced expression of SLC16A1	(Pullen et al., 2011)
miR-31	colorectal cancer	cell-cycle deregulation; reduced apoptosis rate	(Wang et al., 2010b; Cekaite et al., 2012)
miR-320	colon cancer	increased expression of SOX4; inhibition of p53 mediated apoptosis; reduced expression of FOXM1 and FOXQ1; cell-cycle deregulation	(Wan et al., 2015; Vishnubalaji et al., 2016)
miR-338-3p	p53 mutant colorectal cancer	reduced expression of mTOR; increased autophagy and reduced apoptosis rate	(Han et al., 2017)
miR-34a	colon cancer	increased expression of anti-apoptotic protein Bcl-2	(Wang et al., 2010a)
miR-365	colon cancer	increased expression of anti-apoptotic protein Bcl-2	(Nie et al., 2012)

Non-coding RNAs and anticancer drugs

miR-374b	pancreatic cancer	increased ATP7A expression	(Schreiber et al., 2016)
miR-378	liver cancer	increased expression of CYP2E1	(Mohri et al., 2010)
miR-409-3p	colon cancer	increased expression of Beclin-1; increased autophagy pathway	(Tan et al., 2016)
miR-451	colon cancer	increasing the self-renewal of cancer stem cells; increased expression of ABCB1	(Bitarte et al., 2011)
miR-494	colon cancer	reduced expression of dihydropyrimidine dehydrogenase	(Chai et al., 2015)
miR-503-5p	colorectal cancer	reduced expression of apoptotic protein PUMA	(Xu et al., 2017a)
miR-508-5p	gastric cancer	increased expression of ABCB1; increased expression of transcription factor ZNRD1	(Shang et al., 2014)
miR-519d	liver cancer	reduced expression of G1-checkpoint CDK inhibitor p21; reduced apoptosis rate	(Fornari et al., 2012)
miR-522	colon cancer	increased expression of ABCB5	(Yang et al., 2015)
miR-92b	colon cancer	reduced expression of SLC15A and SLC15A1	(Dalmasso et al., 2011)
miR-939	gastric cancer	increased expression of SLC34A2; activation of Ras/MEK/ERK pathway	(Zhang et al., 2017)
miR-96	colorectal cancer	reduced expression of anti-apoptotic proteins XIAP and UBE2N	(Kim et al., 2015)
svRNAb	all GI tumours	reduced expression of CYP3A4	(Persson et al., 2009)
vRNA hvg-1	all GI tumours	transporting drugs away from the target and drug sequestration	(Mossink et al., 2003;Gopinath et al., 2010)
vRNA hvg-2	all GI tumours	transporting drugs away from the target and drug sequestration	(Mossink et al., 2003;Gopinath et al., 2010)

883

884

885 **Table 2: Overview about the different categories of non-coding RNA molecules.**

886

Name	Biological role
circular RNA (circRNA)	involved in forming RNA-protein complex that regulate gene transcription; involved in regulating gene expression at post-transcriptional level by acting as miRNA sponge
endogenous small interfering RNA (endo-siRNA)	involved in repression of transposable elements, chromatin organisation as well as gene regulation at transcriptional and post-transcriptional level
extracellular RNA (exRNA)	involved in intercellular communication and cell regulation
long intergenic non-coding RNA (lincRNA)	involved in gene expression <i>via</i> directing chromatin-modification complexes to specific target regions; lincRNAs located in the cytoplasm function as scaffold to bring together proteins and other RNA categories (especially mRNAs and miRNAs)
long non-coding RNA (lncRNA)	involved in regulation of gene expression <i>via</i> binding to chromatin regulatory proteins; involved in regulating gene expression at post-transcriptional level by acting as microRNA decoys; some lncRNAs are processed into microRNAs
microRNA	involved in fine tuning cell homeostasis by controlling gene expression at post-transcriptional level
miRNA-offset-RNA (moRNA)	unknown
piwi-interacting RNA (piRNA)	involved in maintain germline integrity by repressing transposable elements; involved in mRNA de-adenylation;
ribosomal RNA (rRNA)	component of the ribosomes; involved in protein synthesis
small Cajal body RNA (scaRNA)	component of the Cajal bodies; involved in the biogenesis of small nuclear ribonucleoproteins and by this influence splicing of pre-mRNAs
small interfering RNA (siRNA)	involved in RNA interference pathway as part of anti-viral defence
small nuclear RNA (snRNA)	component of the spliceosome; involved in splicing of pre-mRNAs during post-transcriptional modifications
small nucleolar RNA (snoRNA)	component of the Cajal bodies; involved in modification and processing of snRNA, rRNA and tRNA precursors as well as in mRNA editing
sno-derived RNA (sdRNA)	component of the Cajal bodies; involved in alternative splicing of mRNAs; some sdRNAs control gene expression at post-transcriptional level
transcription initiation RNA (tiRNA)	involved in regulation of RNA polymerase II dependent transcription
transfer RNA (tRNA)	involved in transporting amino acids to the ribosomes during translation
vault RNA (vRNA)	component of the vaults (large ribonucleoprotein complexes in

cytoplasm); unknown function

Table 3: Approved targeted therapies for GI cancer

Abbreviation used: HER2=human epidermal growth factor receptor 2 ; VEGFR=vascular endothelial growth factor receptor; PD-1=programmed cell death protein-1 ; RAF=rapidly accelerated fibrosarcoma; PDGFR=platelet-derived growth factor receptor; c-KIT=SCFR=mast/stem cell growth factor receptor; EGFR=epidermal growth factor receptor; VEGF=vascular endothelial growth factor; RET=rearranged during transfection; MSI-H=microsatellite instability-high

GI cancer	Drug	Target
Gastric cancer	Trastuzumab	HER2
	Ramucirumab	VEGFR-2
	Pembrolizumab	PD-1
Hepatocellular cancer	Sorafenib	RAF, VEGFR-2, VEGFR-3, PDGFR, c-KIT
Colon cancer	Cetuximab, Panitumumab	EGFR
	Bevacizumab	VEGF
	Regorafenib	VEGFR-1, VEGFR-2, VEGFR-3, BRAF, c-KIT, RET, PDGFR
Colon cancer with MSI-H	Pembrolizumab	PD-1

899 **References:**

- 900 Abukhdeir, A.M., and Park, B.H. (2008). P21 and p27: roles in carcinogenesis and
901 drug resistance. *Expert Rev Mol Med* 10, e19.
- 902 Acunzo, M., Romano, G., Wernicke, D., and Croce, C.M. (2015). MicroRNA and
903 cancer--a brief overview. *Adv Biol Regul* 57, 1-9.
- 904 Adams, B.D., Parsons, C., Walker, L., Zhang, W.C., and Slack, F.J. (2017).
905 Targeting noncoding RNAs in disease. *J Clin Invest* 127, 761-771.
- 906 Akita, H., Marquardt, J.U., Durkin, M.E., Kitade, M., Seo, D., Conner, E.A.,
907 Andersen, J.B., Factor, V.M., and Thorgeirsson, S.S. (2014). MYC activates
908 stem-like cell potential in hepatocarcinoma by a p53-dependent mechanism.
909 *Cancer Res* 74, 5903-5913.
- 910 Amankwatia, E.B., Chakravarty, P., Carey, F.A., Weidlich, S., Steele, R.J., Munro,
911 A.J., Wolf, C.R., and Smith, G. (2015). MicroRNA-224 is associated with
912 colorectal cancer progression and response to 5-fluorouracil-based
913 chemotherapy by KRAS-dependent and -independent mechanisms. *Br J*
914 *Cancer* 112, 1480-1490.
- 915 An, X., Sarmiento, C., Tan, T., and Zhu, H. (2017). Regulation of multidrug
916 resistance by microRNAs in anti-cancer therapy. *Acta Pharmaceutica Sinica B*
917 7, 38-51.
- 918 An, Y., Furber, K.L., and Ji, S. (2016). Pseudogenes regulate parental gene
919 expression via ceRNA network. *J Cell Mol Med*.
- 920 Azlan, A., Dzaki, N., and Azzam, G. (2016). Argonaute: The executor of small RNA
921 function. *Journal of Genetics and Genomics* 43, 481-494.
- 922 Balatti, V., Pekarky, Y., and Croce, C.M. (2015). Role of microRNA in chronic
923 lymphocytic leukemia onset and progression. *J Hematol Oncol* 8, 12.
- 924 Bedi, U., Mishra, V.K., Wasilewski, D., Scheel, C., and Johnsen, S.A. (2014).
925 Epigenetic plasticity: a central regulator of epithelial-to-mesenchymal
926 transition in cancer. *Oncotarget* 5, 2016-2029.
- 927 Beermann, J., Piccoli, M.T., Viereck, J., and Thum, T. (2016). Non-coding RNAs in
928 Development and Disease: Background, Mechanisms, and Therapeutic
929 Approaches. *Physiol Rev* 96, 1297-1325.
- 930 Bhutia, Y.D., Hung, S.W., Krentz, M., Patel, D., Lovin, D., Manoharan, R., Thomson,
931 J.M., and Govindarajan, R. (2013). Differential Processing of let-7a
932 Precursors Influences RRM2 Expression and Chemosensitivity in Pancreatic
933 Cancer: Role of LIN-28 and SET Oncoprotein. *Plos One* 8.
- 934 Bian, Z., Jin, L., Zhang, J., Yin, Y., Quan, C., Hu, Y., Feng, Y., Liu, H., Fei, B., Mao,
935 Y., Zhou, L., Qi, X., Huang, S., Hua, D., Xing, C., and Huang, Z. (2016).
936 LncRNA-UCA1 enhances cell proliferation and 5-fluorouracil resistance in
937 colorectal cancer by inhibiting miR-204-5p. *Sci Rep* 6, 23892.
- 938 Bitarte, N., Bandres, E., Boni, V., Zarate, R., Rodriguez, J., Gonzalez-Huarriz, M.,
939 Lopez, I., Javier Sola, J., Alonso, M.M., Fortes, P., and Garcia-Foncillas, J.
940 (2011). MicroRNA-451 is involved in the self-renewal, tumorigenicity, and
941 chemoresistance of colorectal cancer stem cells. *Stem Cells* 29, 1661-1671.
- 942 Blanke, C.D., Demetri, G.D., Von Mehren, M., Heinrich, M.C., Eisenberg, B.,
943 Fletcher, J.A., Corless, C.L., Fletcher, C.D., Roberts, P.J., Heinz, D., Wehre,
944 E., Nikolova, Z., and Joensuu, H. (2008). Long-term results from a
945 randomized phase II trial of standard- versus higher-dose imatinib mesylate

- for patients with unresectable or metastatic gastrointestinal stromal tumors expressing KIT. *J Clin Oncol* 26, 620-625.
- Boni, V., Bitarte, N., Cristobal, I., Zarate, R., Rodriguez, J., Maiello, E., Garcia-Foncillas, J., and Bandres, E. (2010). miR-192/miR-215 Influence 5-Fluorouracil Resistance through Cell Cycle-Mediated Mechanisms Complementary to Its Post-transcriptional Thymidilate Synthase Regulation. *Molecular Cancer Therapeutics* 9, 2265-2275.
- Bottai, G., Pasculli, B., Calin, G.A., and Santarpia, L. (2014). Targeting the microRNA-regulating DNA damage/repair pathways in cancer. *Expert Opinion on Biological Therapy* 14, 1667-1683.
- Braconi, C., Valeri, N., Gasparini, P., Huang, N., Taccioli, C., Nuovo, G., Suzuki, T., Croce, C.M., and Patel, T. (2010). Hepatitis C virus proteins modulate microRNA expression and chemosensitivity in malignant hepatocytes. *Clin Cancer Res* 16, 957-966.
- Bu, P., Chen, K.Y., Chen, J.H., Wang, L., Walters, J., Shin, Y.J., Goerger, J.P., Sun, J., Witherspoon, M., Rakhilin, N., Li, J., Yang, H., Milsom, J., Lee, S., Zipfel, W., Jin, M.M., Gumus, Z.H., Lipkin, S.M., and Shen, X. (2013). A microRNA miR-34a-regulated bimodal switch targets Notch in colon cancer stem cells. *Cell Stem Cell* 12, 602-615.
- Bunz, F., Dutriaux, A., Lengauer, C., Waldman, T., Zhou, S., Brown, J.P., Sedivy, J.M., Kinzler, K.W., and Vogelstein, B. (1998). Requirement for p53 and p21 to sustain G(2) arrest after DNA damage. *Science* 282, 1497-1501.
- Carotenuto, P., Zitto, D., Previdi, M.C., Raj, M., Fassan, M., Lampis, A., Scalafani, F., Lanese, A., Said-Huntingford, I., Hahne, J.C., Young, K., Begum, R., Ethiar, Z., Wotherspoon, A., Starling, N., Sadanadam, A., Cunningham, D., Chau, I., Workman, P., Chopra, R., Valeri, N., and Braconi, C. (2018). MiR1307 mediates pancreatic cancer resistance to FOLFIRINOX chemotherapy by affecting response to DNA damage. *in press*.
- Cekaite, L., Rantala, J.K., Bruun, J., Guriby, M., Agesen, T.H., Danielsen, S.A., Lind, G.E., Nesbakken, A., Kallioniemi, O., Lothe, R.A., and Skotheim, R.I. (2012). MiR-9, -31, and -182 deregulation promote proliferation and tumor cell survival in colon cancer. *Neoplasia* 14, 868-879.
- Chai, J., Dong, W., Xie, C., Wang, L., Han, D.L., Wang, S., Guo, H.L., and Zhang, Z.L. (2015). MicroRNA-494 Sensitizes Colon Cancer Cells to Fluorouracil Through Regulation of DPYD. *Iubmb Life* 67, 191-201.
- Chan, B.A., Jang, R.W., Wong, R.K., Swallow, C.J., Darling, G.E., and Elimova, E. (2016). Improving Outcomes in Resectable Gastric Cancer: A Review of Current and Future Strategies. *Oncology (Williston Park)* 30, 635-645.
- Chapman, P.B., Hauschild, A., Robert, C., Haanen, J.B., Ascierto, P., Larkin, J., Dummer, R., Garbe, C., Testori, A., Maio, M., Hogg, D., Lorigan, P., Lebbe, C., Jouary, T., Schadendorf, D., Ribas, A., O'day, S.J., Sosman, J.A., Kirkwood, J.M., Eggermont, A.M., Dreno, B., Nolop, K., Li, J., Nelson, B., Hou, J., Lee, R.J., Flaherty, K.T., Mcarthur, G.A., and Group, B.-S. (2011). Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 364, 2507-2516.
- Chen, F.J., Sun, M., Li, S.Q., Wu, Q.Q., Ji, L., Liu, Z.L., Zhou, G.Z., Cao, G., Jin, L., Xie, H.W., Wang, C.M., Lv, J., De, W., Wu, M., and Cao, X.F. (2013). Upregulation of the long non-coding RNA HOTAIR promotes esophageal

- squamous cell carcinoma metastasis and poor prognosis. *Mol Carcinog* 52, 908-915.
- Chen, P., Wan, D., Zheng, D., Zheng, Q., Wu, F., and Zhi, Q. (2016). Long non-coding RNA UCA1 promotes the tumorigenesis in pancreatic cancer. *Biomed Pharmacother* 83, 1220-1226.
- Chen, S., Zhu, J., Wang, F., Guan, Z., Ge, Y., Yang, X., and Cai, J. (2017). LncRNAs and their role in cancer stem cells. *Oncotarget* 8, 110685-110692.
- Cheng, A.S., Lau, S.S., Chen, Y., Kondo, Y., Li, M.S., Feng, H., Ching, A.K., Cheung, K.F., Wong, H.K., Tong, J.H., Jin, H., Choy, K.W., Yu, J., To, K.F., Wong, N., Huang, T.H., and Sung, J.J. (2011). EZH2-mediated concordant repression of Wnt antagonists promotes beta-catenin-dependent hepatocarcinogenesis. *Cancer Res* 71, 4028-4039.
- Choudhuri, S. (2010). Small Noncoding RNAs: Biogenesis, Function, and Emerging Significance in Toxicology. *Journal of Biochemical and Molecular Toxicology* 24, 195-216.
- Chowdhury, D., Choi, Y.E., and Brault, M.E. (2013). DNA DAMAGE - OPINION Charity begins at home: non-coding RNA functions in DNA repair. *Nature Reviews Molecular Cell Biology* 14, 181-189.
- Choy, K.R., and Watters, D.J. (2018). Neurodegeneration in ataxia-telangiectasia: Multiple roles of ATM kinase in cellular homeostasis. *Dev Dyn* 247, 33-46.
- Claycomb, J.M. (2014). Ancient endo-siRNA pathways reveal new tricks. *Curr Biol* 24, R703-715.
- Colas, C., Ung, P.M.U., and Schlessinger, A. (2016). SLC transporters: structure, function, and drug discovery. *Medchemcomm* 7, 1069-1081.
- Cortez, M.A., Ivan, C., Valdecanas, D., Wang, X., Peltier, H.J., Ye, Y., Araujo, L., Carbone, D.P., Shilo, K., Giri, D.K., Kelnar, K., Martin, D., Komaki, R., Gomez, D.R., Krishnan, S., Calin, G.A., Bader, A.G., and Welsh, J.W. (2016). PDL1 Regulation by p53 via miR-34. *J Natl Cancer Inst* 108.
- Cox, J., and Weinman, S. (2016). Mechanisms of doxorubicin resistance in hepatocellular carcinoma. *Hepat Oncol* 3, 57-59.
- Dalmasso, G., Hang, T.T.N., Yan, Y.T., Laroui, H., Charania, M.A., Obertone, T.S., Sitaraman, S.V., and Merlin, D. (2011). MicroRNA-92b regulates expression of the oligopeptide transporter PepT1 in intestinal epithelial cells. *American Journal of Physiology-Gastrointestinal and Liver Physiology* 300, G52-G59.
- De Almeida, R.A., Fraczek, M.G., Parker, S., Delneri, D., and O'keefe, R.T. (2016). Non-coding RNAs and disease: the classical ncRNAs make a comeback. *Biochem Soc Trans* 44, 1073-1078.
- Dean, M., Hamon, Y., and Chimini, G. (2001). The human ATP-binding cassette (ABC) transporter superfamily. *J Lipid Res* 42, 1007-1017.
- Deng, Y., Huang, Z., Xu, Y., Jin, J., Zhuo, W., Zhang, C., Zhang, X., Shen, M., Yan, X., Wang, L., Wang, X., Kang, Y., Si, J., and Zhou, T. (2014). MiR-215 modulates gastric cancer cell proliferation by targeting RB1. *Cancer Lett* 342, 27-35.
- Ding, C., Cheng, S., Yang, Z., Lv, Z., Xiao, H., Du, C., Peng, C., Xie, H., Zhou, L., Wu, J., and Zheng, S. (2014). Long non-coding RNA HOTAIR promotes cell migration and invasion via down-regulation of RNA binding motif protein 38 in hepatocellular carcinoma cells. *Int J Mol Sci* 15, 4060-4076.

- Donahue, T.R., Nguyen, A.H., Moughan, J., Li, L., Tatishchev, S., Toste, P., and Farrell, J.J. (2014). Stromal microRNA-21 levels predict response to 5-fluorouracil in patients with pancreatic cancer. *J Surg Oncol* 110, 952-959.
- Dong, J., Zhao, Y.P., Zhou, L., Zhang, T.P., and Chen, G. (2011). Bcl-2 upregulation induced by miR-21 via a direct interaction is associated with apoptosis and chemoresistance in MIA PaCa-2 pancreatic cancer cells. *Arch Med Res* 42, 8-14.
- Dong, R., Liu, G.B., Liu, B.H., Chen, G., Li, K., Zheng, S., and Dong, K.R. (2016). Targeting long non-coding RNA-TUG1 inhibits tumor growth and angiogenesis in hepatoblastoma. *Cell Death & Disease* 7.
- Douillard, J.Y., Rong, A., and Sidhu, R. (2013). RAS mutations in colorectal cancer. *N Engl J Med* 369, 2159-2160.
- Du, B., and Shim, J.S. (2016). Targeting Epithelial-Mesenchymal Transition (EMT) to Overcome Drug Resistance in Cancer. *Molecules* 21.
- Eddy, S.R. (2001). Non-coding RNA genes and the modern RNA world. *Nature Reviews Genetics* 2, 919-929.
- Edelbrock, M.A., Kaliyaperumal, S., and Williams, K.J. (2013). Structural, molecular and cellular functions of MSH2 and MSH6 during DNA mismatch repair, damage signaling and other noncanonical activities. *Mutat Res* 743-744, 53-66.
- El-Deiry, W.S., Tokino, T., Velculescu, V.E., Levy, D.B., Parsons, R., Trent, J.M., Lin, D., Mercer, W.E., Kinzler, K.W., and Vogelstein, B. (1993). WAF1, a potential mediator of p53 tumor suppression. *Cell* 75, 817-825.
- Endo, H., Shiroki, T., Nakagawa, T., Yokoyama, M., Tamai, K., Yamanami, H., Fujiya, T., Sato, I., Yamaguchi, K., Tanaka, N., Iijima, K., Shimosegawa, T., Sugamura, K., and Satoh, K. (2013). Enhanced expression of long non-coding RNA HOTAIR is associated with the development of gastric cancer. *PLoS One* 8, e77070.
- Esteller, M. (2011). Non-coding RNAs in human disease. *Nat Rev Genet* 12, 861-874.
- Evans, J., Essex, A., Xin, H., Amitai, N., Brinton, L., Griner, E., Reproducibility Project: Cancer, B., and Reproducibility Project Cancer, B. (2015). Registered report: Wnt activity defines colon cancer stem cells and is regulated by the microenvironment. *Elife* 4.
- Evans, J.R., Feng, F.Y., and Chinnaiyan, A.M. (2016). The bright side of dark matter: lncRNAs in cancer. *J Clin Invest* 126, 2775-2782.
- Fabbri, M., Valeri, N., and Calin, G.A. (2009). MicroRNAs and genomic variations: from Proteus tricks to Prometheus gift. *Carcinogenesis* 30, 912-917.
- Fan, Y., Dutta, J., Gupta, N., Fan, G., and Gelinas, C. (2008). Regulation of programmed cell death by NF-kappaB and its role in tumorigenesis and therapy. *Adv Exp Med Biol* 615, 223-250.
- Fan, Y., Shen, B., Tan, M., Mu, X., Qin, Y., Zhang, F., and Liu, Y. (2014a). Long non-coding RNA UCA1 increases chemoresistance of bladder cancer cells by regulating Wnt signaling. *FEBS J* 281, 1750-1758.
- Fan, Y., Shen, B., Tan, M., Mu, X., Qin, Y., Zhang, F., and Liu, Y. (2014b). TGF-beta-induced upregulation of malat1 promotes bladder cancer metastasis by associating with suz12. *Clin Cancer Res* 20, 1531-1541.

- Fang, L., Li, H., Wang, L., Hu, J., Jin, T., Wang, J., and Yang, B.B. (2014). MicroRNA-17-5p promotes chemotherapeutic drug resistance and tumour metastasis of colorectal cancer by repressing PTEN expression. *Oncotarget* 5, 2974-2987.
- Fang, L., Zhang, J., Zhang, H., Yang, X., Jin, X., Zhang, L., Skalnik, D.G., Jin, Y., Zhang, Y., Huang, X., Li, J., and Wong, J. (2016a). H3K4 Methyltransferase Set1a Is A Key Oct4 Coactivator Essential for Generation of Oct4 Positive Inner Cell Mass. *Stem Cells* 34, 565-580.
- Fang, Q., Chen, X., and Zhi, X. (2016b). Long Non-Coding RNA (LncRNA) Urothelial Carcinoma Associated 1 (UCA1) Increases Multi-Drug Resistance of Gastric Cancer via Downregulating miR-27b. *Med Sci Monit* 22, 3506-3513.
- Fang, X.Y., Pan, H.F., Leng, R.X., and Ye, D.Q. (2015). Long noncoding RNAs: novel insights into gastric cancer. *Cancer Lett* 356, 357-366.
- Fornari, F., Gramantieri, L., Giovannini, C., Veronese, A., Ferracin, M., Sabbioni, S., Calin, G.A., Grazi, G.L., Croce, C.M., Tavalari, S., Chieco, P., Negrini, M., and Bolondi, L. (2009). MiR-122/cyclin G1 interaction modulates p53 activity and affects doxorubicin sensitivity of human hepatocarcinoma cells. *Cancer Res* 69, 5761-5767.
- Fornari, F., Milazzo, M., Chieco, P., Negrini, M., Calin, G.A., Grazi, G.L., Pollutri, D., Croce, C.M., Bolondi, L., and Gramantieri, L. (2010). MiR-199a-3p regulates mTOR and c-Met to influence the doxorubicin sensitivity of human hepatocarcinoma cells. *Cancer Res* 70, 5184-5193.
- Fornari, F., Milazzo, M., Chieco, P., Negrini, M., Marasco, E., Capranico, G., Mantovani, V., Marinello, J., Sabbioni, S., Callegari, E., Cescon, M., Ravaioli, M., Croce, C.M., Bolondi, L., and Gramantieri, L. (2012). In hepatocellular carcinoma miR-519d is up-regulated by p53 and DNA hypomethylation and targets CDKN1A/p21, PTEN, AKT3 and TIMP2. *J Pathol* 227, 275-285.
- Frampton, A.E., Castellano, L., Colombo, T., Giovannetti, E., Krell, J., Jacob, J., Pellegrino, L., Roca-Alonso, L., Funel, N., Gall, T.M., De Giorgio, A., Pinho, F.G., Fulci, V., Britton, D.J., Ahmad, R., Habib, N.A., Coombes, R.C., Harding, V., Knosel, T., Stebbing, J., and Jiao, L.R. (2014a). MicroRNAs cooperatively inhibit a network of tumor suppressor genes to promote pancreatic tumor growth and progression. *Gastroenterology* 146, 268-277 e218.
- Frampton, A.E., Giovannetti, E., Jamieson, N.B., Krell, J., Gall, T.M., Stebbing, J., Jiao, L.R., and Castellano, L. (2014b). A microRNA meta-signature for pancreatic ductal adenocarcinoma. *Expert Rev Mol Diagn* 14, 267-271.
- Gagez, A.L., Duroux-Richard, I., Lepretre, S., Orsini-Piocelle, F., Letestu, R., De Guibert, S., Tuailon, E., Leblond, V., Khalifa, O., Gouilleux-Gruart, V., Banos, A., Tournilhac, O., Dupuis, J., Jorgensen, C., Cartron, G., and Apparailly, F. (2017). miR-125b and miR-532-3p predict the efficiency of rituximab-mediated lymphodepletion in chronic lymphocytic leukemia patients. A French Innovative Leukemia Organization study. *Haematologica* 102, 746-754.
- Gardiner, A.S., Twiss, J.L., and Perrone-Bizzozero, N.I. (2015). Competing Interactions of RNA-Binding Proteins, MicroRNAs, and Their Targets Control Neuronal Development and Function. *Biomolecules* 5, 2903-2918.
- Garg, M. (2015). Emerging role of microRNAs in cancer stem cells: Implications in cancer therapy. *World J Stem Cells* 7, 1078-1089.

- Ge, X.S., Ma, H.J., Zheng, X.H., Ruan, H.L., Liao, X.Y., Xue, W.Q., Chen, Y.B., Zhang, Y., and Jia, W.H. (2013). HOTAIR, a prognostic factor in esophageal squamous cell carcinoma, inhibits WIF-1 expression and activates Wnt pathway. *Cancer Sci* 104, 1675-1682.
- Geiger, J., and Dalgaard, L.T. (2016). Interplay of mitochondrial metabolism and microRNAs. *Cell Mol Life Sci*.
- Geisler, S., and Collier, J. (2013). RNA in unexpected places: long non-coding RNA functions in diverse cellular contexts. *Nat Rev Mol Cell Biol* 14, 699-712.
- Geng, Y.J., Xie, S.L., Li, Q., Ma, J., and Wang, G.Y. (2011). Large Intervening Non-coding RNA HOTAIR is Associated with Hepatocellular Carcinoma Progression. *Journal of International Medical Research* 39, 2119-2128.
- Giovannetti, E., Funel, N., Peters, G.J., Del Chiaro, M., Erozcenci, L.A., Vasile, E., Leon, L.G., Pollina, L.E., Groen, A., Falcone, A., Danesi, R., Campani, D., Verheul, H.M., and Boggi, U. (2010). MicroRNA-21 in pancreatic cancer: correlation with clinical outcome and pharmacologic aspects underlying its role in the modulation of gemcitabine activity. *Cancer Res* 70, 4528-4538.
- Gong, J., Cui, Z., Li, L., Ma, Q., Wang, Q., Gao, Y., and Sun, H. (2015). MicroRNA-25 promotes gastric cancer proliferation, invasion, and migration by directly targeting F-box and WD-40 Domain Protein 7, FBXW7. *Tumour Biol* 36, 7831-7840.
- Gonzalez, C.D., Alvarez, S., Ropolo, A., Rosenzvit, C., Bagnes, M.F., and Vaccaro, M.I. (2014). Autophagy, Warburg, and Warburg reverse effects in human cancer. *Biomed Res Int* 2014, 926729.
- Gopinath, S.C., Wadhwa, R., and Kumar, P.K. (2010). Expression of noncoding vault RNA in human malignant cells and its importance in mitoxantrone resistance. *Mol Cancer Res* 8, 1536-1546.
- Gottesman, M.M., Fojo, T., and Bates, S.E. (2002). Multidrug resistance in cancer: Role of ATP-dependent transporters. *Nature Reviews Cancer* 2, 48-58.
- Gozuacik, D., Akkoc, Y., Ozturk, D.G., and Kocak, M. (2017). Autophagy-Regulating microRNAs and Cancer. *Front Oncol* 7, 65.
- Granados-Riveron, J.T., and Aquino-Jarquin, G. (2016). The complexity of the translation ability of circRNAs. *Biochim Biophys Acta* 1859, 1245-1251.
- Grasedieck, S., Scholer, N., Bommer, M., Niess, J.H., Tumani, H., Rouhi, A., Bloehdorn, J., Liebisch, P., Mertens, D., Dohner, H., Buske, C., Langer, C., and Kuchenbauer, F. (2012). Impact of serum storage conditions on microRNA stability. *Leukemia* 26, 2414-2416.
- Graziano, F., Canestrari, E., Loupakis, F., Ruzzo, A., Galluccio, N., Santini, D., Rocchi, M., Vincenzi, B., Salvatore, L., Cremolini, C., Spoto, C., Catalano, V., D'emidio, S., Giordani, P., Tonini, G., Falcone, A., and Magnani, M. (2010). Genetic modulation of the Let-7 microRNA binding to KRAS 3'-untranslated region and survival of metastatic colorectal cancer patients treated with salvage cetuximab-irinotecan. *Pharmacogenomics J* 10, 458-464.
- Grothey, A., Van Cutsem, E., Sobrero, A., Siena, S., Falcone, A., Ychou, M., Humblet, Y., Bouche, O., Mineur, L., Barone, C., Adenis, A., Tabernero, J., Yoshino, T., Lenz, H.J., Goldberg, R.M., Sargent, D.J., Cihon, F., Cupit, L., Wagner, A., Laurent, D., and Group, C.S. (2013). Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an

- international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 381, 303-312.
- Guengerich, F.P., and Shimada, T. (1991). Oxidation of toxic and carcinogenic chemicals by human cytochrome P-450 enzymes. *Chem Res Toxicol* 4, 391-407.
- Gui, X., Li, H., Li, T., Pu, H., and Lu, D. (2015). Long Noncoding RNA CUDR Regulates HULC and beta-Catenin to Govern Human Liver Stem Cell Malignant Differentiation. *Mol Ther* 23, 1843-1853.
- Guo, J.U., Agarwal, V., Guo, H., and Bartel, D.P. (2014). Expanded identification and characterization of mammalian circular RNAs. *Genome Biol* 15, 409.
- Gupta, R.A., Shah, N., Wang, K.C., Kim, J., Horlings, H.M., Wong, D.J., Tsai, M.C., Hung, T., Argani, P., Rinn, J.L., Wang, Y., Brzoska, P., Kong, B., Li, R., West, R.B., Van De Vijver, M.J., Sukumar, S., and Chang, H.Y. (2010). Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* 464, 1071-1076.
- Guttman, M., Amit, I., Garber, M., French, C., Lin, M.F., Feldser, D., Huarte, M., Zuk, O., Carey, B.W., Cassady, J.P., Cabili, M.N., Jaenisch, R., Mikkelsen, T.S., Jacks, T., Hacohen, N., Bernstein, B.E., Kellis, M., Regev, A., Rinn, J.L., and Lander, E.S. (2009). Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* 458, 223-227.
- Guttman, M., Donaghey, J., Carey, B.W., Garber, M., Grenier, J.K., Munson, G., Young, G., Lucas, A.B., Ach, R., Bruhn, L., Yang, X.P., Amit, I., Meissner, A., Regev, A., Rinn, J.L., Root, D.E., and Lander, E.S. (2011). lincRNAs act in the circuitry controlling pluripotency and differentiation. *Nature* 477, 295-U260.
- Ha, T.Y. (2011a). MicroRNAs in Human Diseases: From Cancer to Cardiovascular Disease. *Immune Netw* 11, 135-154.
- Ha, T.Y. (2011b). MicroRNAs in Human Diseases: From Lung, Liver and Kidney Diseases to Infectious Disease, Sickle Cell Disease and Endometrium Disease. *Immune Netw* 11, 309-323.
- Ha, T.Y. (2011c). The Role of MicroRNAs in Regulatory T Cells and in the Immune Response. *Immune Netw* 11, 11-41.
- Haenisch, S., Werk, A.N., and Cascorbi, I. (2014). MicroRNAs and their relevance to ABC transporters. *British Journal of Clinical Pharmacology* 77, 587-596.
- Han, J., Li, J., Tang, K., Zhang, H., Guo, B., Hou, N., and Huang, C. (2017). miR-338-3p confers 5-fluorouracil resistance in p53 mutant colon cancer cells by targeting the mammalian target of rapamycin. *Exp Cell Res* 360, 328-336.
- Han, P., and Chang, C.P. (2015). Long non-coding RNA and chromatin remodeling. *RNA Biol* 12, 1094-1098.
- Han, Y., Yang, Y.N., Yuan, H.H., Zhang, T.T., Sui, H., Wei, X.L., Liu, L., Huang, P., Zhang, W.J., and Bai, Y.X. (2014a). UCA1, a long non-coding RNA up-regulated in colorectal cancer influences cell proliferation, apoptosis and cell cycle distribution. *Pathology* 46, 396-401.
- Han, Y., Ye, J., Wu, D., Wu, P., Chen, Z., Chen, J., Gao, S., and Huang, J. (2014b). LEIGC long non-coding RNA acts as a tumor suppressor in gastric carcinoma by inhibiting the epithelial-to-mesenchymal transition. *BMC Cancer* 14, 932.
- Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: the next generation. *Cell* 144, 646-674.

- 1229 Hang, Q., Sun, R., Jiang, C., and Li, Y. (2015). Notch 1 promotes cisplatin-resistant
1230 gastric cancer formation by upregulating lncRNA AK022798 expression.
1231 *Anticancer Drugs* 26, 632-640.
- 1232 Hansen, T.F., Carlsen, A.L., Heegaard, N.H., Sorensen, F.B., and Jakobsen, A.
1233 (2015). Changes in circulating microRNA-126 during treatment with
1234 chemotherapy and bevacizumab predicts treatment response in patients with
1235 metastatic colorectal cancer. *Br J Cancer* 112, 624-629.
- 1236 Harper, J.W., and Elledge, S.J. (2007). The DNA damage response: Ten years after.
1237 *Molecular Cell* 28, 739-745.
- 1238 Hasegawa, S., Eguchi, H., Nagano, H., Konno, M., Tomimaru, Y., Wada, H., Hama,
1239 N., Kawamoto, K., Kobayashi, S., Nishida, N., Koseki, J., Nishimura, T.,
1240 Gotoh, N., Ohno, S., Yabuta, N., Nojima, H., Mori, M., Doki, Y., and Ishii, H.
1241 (2014). MicroRNA-1246 expression associated with CCNG2-mediated
1242 chemoresistance and stemness in pancreatic cancer. *Br J Cancer* 111, 1572-
1243 1580.
- 1244 He, H., Tian, W., Chen, H., and Deng, Y. (2016). MicroRNA-101 sensitizes
1245 hepatocellular carcinoma cells to doxorubicin-induced apoptosis via targeting
1246 Mcl-1. *Mol Med Rep* 13, 1923-1929.
- 1247 He, J., Qi, H., Chen, F., and Cao, C. (2017). MicroRNA-25 contributes to cisplatin
1248 resistance in gastric cancer cells by inhibiting forkhead box O3a. *Oncol Lett*
1249 14, 6097-6102.
- 1250 He, L., and Hannon, G.J. (2004). MicroRNAs: small RNAs with a big role in gene
1251 regulation. *Nat Rev Genet* 5, 522-531.
- 1252 He, Y., Meng, X.M., Huang, C., Wu, B.M., Zhang, L., Lv, X.W., and Li, J. (2014).
1253 Long noncoding RNAs: Novel insights into hepatocellular carcinoma. *Cancer*
1254 *Lett* 344, 20-27.
- 1255 Heery, R., Finn, S.P., Cuffe, S., and Gray, S.G. (2017). Long Non-Coding RNAs: Key
1256 Regulators of Epithelial-Mesenchymal Transition, Tumour Drug Resistance
1257 and Cancer Stem Cells. *Cancers (Basel)* 9.
- 1258 Helleday, T., Petermann, E., Lundin, C., Hodgson, B., and Sharma, R.A. (2008).
1259 DNA repair pathways as targets for cancer therapy. *Nature Reviews Cancer*
1260 8, 193-204.
- 1261 Herriges, M.J., Swarr, D.T., Morley, M.P., Rath, K.S., Peng, T., Stewart, K.M., and
1262 Morrissey, E.E. (2014). Long noncoding RNAs are spatially correlated with
1263 transcription factors and regulate lung development. *Genes & Development*
1264 28, 1363-1379.
- 1265 Hiyama, E., and Hiyama, K. (2007). Telomere and telomerase in stem cells. *Br J*
1266 *Cancer* 96, 1020-1024.
- 1267 Hoeijmakers, J.H.J. (2001). Genome maintenance mechanisms for preventing
1268 cancer. *Nature* 411, 366-374.
- 1269 Holohan, C., Van Schaeybroeck, S., Longley, D.B., and Johnston, P.G. (2013).
1270 Cancer drug resistance: an evolving paradigm. *Nat Rev Cancer* 13, 714-726.
- 1271 Housman, G., Byler, S., Heerboth, S., Lapinska, K., Longacre, M., Snyder, N., and
1272 Sarkar, S. (2014). Drug resistance in cancer: an overview. *Cancers (Basel)* 6,
1273 1769-1792.
- 1274 Hsu, S.H., Wang, B., Kota, J., Yu, J., Costinean, S., Kutay, H., Yu, L., Bai, S., La
1275 Perle, K., Chivukula, R.R., Mao, H., Wei, M., Clark, K.R., Mendell, J.R.,
1276 Caligiuri, M.A., Jacob, S.T., Mendell, J.T., and Ghoshal, K. (2012). Essential

- 1277 metabolic, anti-inflammatory, and anti-tumorigenic functions of miR-122 in
1278 liver. *J Clin Invest* 122, 2871-2883.
- 1279 Huang, H., and Tindall, D.J. (2006). FOXO factors: a matter of life and death. *Future*
1280 *Oncol* 2, 83-89.
- 1281 Huang, M.D., Chen, W.M., Qi, F.Z., Sun, M., Xu, T.P., Ma, P., and Shu, Y.Q. (2015).
1282 Long non-coding RNA TUG1 is up-regulated in hepatocellular carcinoma and
1283 promotes cell growth and apoptosis by epigenetically silencing of KLF2.
1284 *Molecular Cancer* 14.
- 1285 Huang, X., Xiao, R., Pan, S., Yang, X., Yuan, W., Tu, Z., Xu, M., Zhu, Y., Yin, Q.,
1286 Wu, Y., Hu, W., Shao, L., Xiong, J., and Zhang, Q. (2017). Uncovering the
1287 roles of long non-coding RNAs in cancer stem cells. *J Hematol Oncol* 10, 62.
- 1288 Huang, Z., Zhou, L., Chen, Z., Nice, E.C., and Huang, C. (2016). Stress
1289 management by autophagy: Implications for chemoresistance. *Int J Cancer*
1290 139, 23-32.
- 1291 Hung, A.Y., Canning, C.A., Patel, K.M., Holland, J.M., and Kachnic, L.A. (2006).
1292 Radiation therapy for gastrointestinal cancer. *Hematol Oncol Clin North Am*
1293 20, 287-320.
- 1294 Hung, T., Wang, Y.L., Lin, M.F., Koegel, A.K., Kotake, Y., Grant, G.D., Horlings,
1295 H.M., Shah, N., Umbricht, C., Wang, P., Wang, Y., Kong, B., Langerod, A.,
1296 Borresen-Dale, A.L., Kim, S.K., Van De Vijver, M., Sukumar, S., Whitfield,
1297 M.L., Kellis, M., Xiong, Y., Wong, D.J., and Chang, H.Y. (2011). Extensive
1298 and coordinated transcription of noncoding RNAs within cell-cycle promoters.
1299 *Nature Genetics* 43, 621-U196.
- 1300 Ikemura, K., Iwamoto, T., and Okuda, M. (2014). MicroRNAs as regulators of drug
1301 transporters, drug-metabolizing enzymes, and tight junctions: Implication for
1302 intestinal barrier function. *Pharmacology & Therapeutics* 143, 217-224.
- 1303 Ikemura, K., Yamamoto, M., Miyazaki, S., Mizutani, H., Iwamoto, T., and Okuda, M.
1304 (2013). MicroRNA-145 Post-transcriptionally Regulates the Expression and
1305 Function of P-glycoprotein in Intestinal Epithelial Cells. *Molecular*
1306 *Pharmacology* 83, 399-405.
- 1307 Iorio, M.V., and Croce, C.M. (2012). microRNA involvement in human cancer.
1308 *Carcinogenesis* 33, 1126-1133.
- 1309 Ischenko, I., Seeliger, H., Kleespies, A., Angele, M.K., Eichhorn, M.E., Jauch, K.W.,
1310 and Bruns, C.J. (2010). Pancreatic cancer stem cells: new understanding of
1311 tumorigenesis, clinical implications. *Langenbecks Arch Surg* 395, 1-10.
- 1312 Ismael, H.N., Denbo, J., Cox, S., Crane, C.H., Das, P., Krishnan, S., Schroff, R.T.,
1313 Javle, M., Conrad, C., Vauthey, J., and Aloia, T. (2016). Biologic mesh spacer
1314 placement facilitates safe delivery of dose-intense radiation therapy: A novel
1315 treatment option for unresectable liver tumors. *Eur J Surg Oncol* 42, 1591-
1316 1596.
- 1317 Jackson, S.P., and Bartek, J. (2009). The DNA-damage response in human biology
1318 and disease. *Nature* 461, 1071-1078.
- 1319 Jakhetiya, A., Garg, P.K., Prakash, G., Sharma, J., Pandey, R., and Pandey, D.
1320 (2016). Targeted therapy of gastrointestinal stromal tumours. *World J*
1321 *Gastrointest Surg* 8, 345-352.
- 1322 Janssen, H.L., Reesink, H.W., Lawitz, E.J., Zeuzem, S., Rodriguez-Torres, M., Patel,
1323 K., Van Der Meer, A.J., Patick, A.K., Chen, A., Zhou, Y., Persson, R., King,

- B.D., Kauppinen, S., Levin, A.A., and Hodges, M.R. (2013). Treatment of HCV infection by targeting microRNA. *N Engl J Med* 368, 1685-1694.
- Jiang, L., Wang, W.C., Li, G.L., Sun, C.L., Ren, Z.Q., Sheng, H.H., Gao, H.J., Wang, C.F., and Yu, H. (2016). High TUG1 expression is associated with chemotherapy resistance and poor prognosis in esophageal squamous cell carcinoma. *Cancer Chemotherapy and Pharmacology* 78, 333-339.
- Jiang, M., Huang, O., Xie, Z., Wu, S., Zhang, X., Shen, A., Liu, H., Chen, X., Wu, J., Lou, Y., Mao, Y., Sun, K., Hu, S., Geng, M., and Shen, K. (2014). A novel long non-coding RNA-ARA: adriamycin resistance-associated. *Biochem Pharmacol* 87, 254-283.
- Jiao, C., Song, Z., Chen, J., Zhong, J., Cai, W., Tian, S., Chen, S., Yi, Y., and Xiao, Y. (2016). lncRNA-UCA1 enhances cell proliferation through functioning as a ceRNA of Sox4 in esophageal cancer. *Oncol Rep* 36, 2960-2966.
- Jiao, F., Hu, H., Han, T., Yuan, C., Wang, L., Jin, Z., Guo, Z., and Wang, L. (2015). Long noncoding RNA MALAT-1 enhances stem cell-like phenotypes in pancreatic cancer cells. *Int J Mol Sci* 16, 6677-6693.
- Jiao, F., Hu, H., Yuan, C., Wang, L., Jiang, W., Jin, Z., Guo, Z., and Wang, L. (2014). Elevated expression level of long noncoding RNA MALAT-1 facilitates cell growth, migration and invasion in pancreatic cancer. *Oncol Rep* 32, 2485-2492.
- Jin, W., Chen, L., Cai, X., Zhang, Y., Zhang, J., Ma, D., Cai, X., Fu, T., Yu, Z., Yu, F., and Chen, G. (2017). Long non-coding RNA TUC338 is functionally involved in sorafenib-sensitized hepatocarcinoma cells by targeting RASAL1. *Oncol Rep* 37, 273-280.
- Jonker, D.J., O'callaghan, C.J., Karapetis, C.S., Zalcberg, J.R., Tu, D., Au, H.J., Berry, S.R., Krahn, M., Price, T., Simes, R.J., Tebbutt, N.C., Van Hazel, G., Wierzbicki, R., Langer, C., and Moore, M.J. (2007). Cetuximab for the treatment of colorectal cancer. *N Engl J Med* 357, 2040-2048.
- Kang, Y.K., Boku, N., Satoh, T., Ryu, M.H., Chao, Y., Kato, K., Chung, H.C., Chen, J.S., Muro, K., Kang, W.K., Yeh, K.H., Yoshikawa, T., Oh, S.C., Bai, L.Y., Tamura, T., Lee, K.W., Hamamoto, Y., Kim, J.G., Chin, K., Oh, D.Y., Minashi, K., Cho, J.Y., Tsuda, M., and Chen, L.T. (2017). Nivolumab in patients with advanced gastric or gastro-oesophageal junction cancer refractory to, or intolerant of, at least two previous chemotherapy regimens (ONO-4538-12, ATTRACTION-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 390, 2461-2471.
- Kano, M., Seki, N., Kikkawa, N., Fujimura, L., Hoshino, I., Akutsu, Y., Chiyomaru, T., Enokida, H., Nakagawa, M., and Matsubara, H. (2010). miR-145, miR-133a and miR-133b: tumor-suppressive miRNAs target FSCN1 in esophageal squamous cell carcinoma. *International Journal of Cancer* 127, 2804-2814.
- Karaayvaz, M., Zhai, H., and Ju, J. (2013). miR-129 promotes apoptosis and enhances chemosensitivity to 5-fluorouracil in colorectal cancer. *Cell Death Dis* 4, e659.
- Kathawala, R.J., Gupta, P., Ashby, C.R., Jr., and Chen, Z.S. (2015). The modulation of ABC transporter-mediated multidrug resistance in cancer: a review of the past decade. *Drug Resist Updat* 18, 1-17.
- Khan, K.H., Yap, T.A., Yan, L., and Cunningham, D. (2013). Targeting the PI3K-AKT-mTOR signaling network in cancer. *Chin J Cancer* 32, 253-265.

- Khurana, E., Fu, Y., Chakravarty, D., Demichelis, F., Rubin, M.A., and Gerstein, M. (2016). Role of non-coding sequence variants in cancer. *Nat Rev Genet* 17, 93-108.
- Kim, G. (2017). nab-Paclitaxel for the treatment of pancreatic cancer. *Cancer Manag Res* 9, 85-96.
- Kim, K., Jutooru, I., Chadalapaka, G., Johnson, G., Frank, J., Burghardt, R., Kim, S., and Safe, S. (2013). HOTAIR is a negative prognostic factor and exhibits pro-oncogenic activity in pancreatic cancer. *Oncogene* 32, 1616-1625.
- Kim, S.A., Kim, I., Yoon, S.K., Lee, E.K., and Kuh, H.J. (2015). Indirect modulation of sensitivity to 5-fluorouracil by microRNA-96 in human colorectal cancer cells. *Arch Pharm Res* 38, 239-248.
- King, J., Palmer, D.H., Johnson, P., Ross, P., Hubner, R.A., Sumpter, K., Darby, S., Braconi, C., Iwaji, C., Swinson, D., Collins, P., Patel, K., Nobes, J., Muazzam, I., Blesing, C., Kirkwood, A., Nash, S., and Meyer, T. (2017). Sorafenib for the Treatment of Advanced Hepatocellular Cancer - a UK Audit. *Clin Oncol (R Coll Radiol)* 29, 256-262.
- Kogo, R., Shimamura, T., Mimori, K., Kawahara, K., Imoto, S., Sudo, T., Tanaka, F., Shibata, K., Suzuki, A., Komune, S., Miyano, S., and Mori, M. (2011). Long Noncoding RNA HOTAIR Regulates Polycomb-Dependent Chromatin Modification and Is Associated with Poor Prognosis in Colorectal Cancers. *Cancer Research* 71, 6320-6326.
- Kornienko, A.E., Guenzl, P.M., Barlow, D.P., and Pauler, F.M. (2013). Gene regulation by the act of long non-coding RNA transcription. *BMC Biol* 11, 59.
- Korpanty, G.J., Graham, D.M., Vincent, M.D., and Leighl, N.B. (2014). Biomarkers That Currently Affect Clinical Practice in Lung Cancer: EGFR, ALK, MET, ROS-1, and KRAS. *Front Oncol* 4, 204.
- Koturbash, I., Beland, F.A., and Pogribny, I.P. (2012). Role of microRNAs in the regulation of drug metabolizing and transporting genes and the response to environmental toxicants. *Expert Opinion on Drug Metabolism & Toxicology* 8, 597-606.
- Kwak, E.L., Bang, Y.J., Camidge, D.R., Shaw, A.T., Solomon, B., Maki, R.G., Ou, S.H., Dezube, B.J., Janne, P.A., Costa, D.B., Varella-Garcia, M., Kim, W.H., Lynch, T.J., Fidias, P., Stubbs, H., Engelman, J.A., Sequist, L.V., Tan, W., Gandhi, L., Mino-Kenudson, M., Wei, G.C., Shreeve, S.M., Ratain, M.J., Settleman, J., Christensen, J.G., Haber, D.A., Wilner, K., Salgia, R., Shapiro, G.I., Clark, J.W., and Iafrate, A.J. (2010). Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med* 363, 1693-1703.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., and Tuschl, T. (2002). Identification of tissue-specific microRNAs from mouse. *Curr Biol* 12, 735-739.
- Lan, W.G., Xu, D.H., Xu, C., Ding, C.L., Ning, F.L., Zhou, Y.L., Ma, L.B., Liu, C.M., and Han, X. (2016). Silencing of long non-coding RNA ANRIL inhibits the development of multidrug resistance in gastric cancer cells. *Oncology Reports* 36, 263-270.
- Langenberger, D., Bermudez-Santana, C., Hertel, J., Hoffmann, S., Khaitovich, P., and Stadler, P.F. (2009). Evidence for human microRNA-offset RNAs in small RNA sequencing data. *Bioinformatics* 25, 2298-2301.

- Laurent-Puig, P., Grisoni, M.L., Heinemann, V., Bonnetain, F., Fontaine, K., Vazart, C., Decaulne, V., Rousseau, F., Courtieu, B., Libaert, F., Jung, A., Neureiter, D., Thiebaut, R., and Stinzing, S. (2016). MiR 31 3p as a predictive biomarker of cetuximab efficacy effect in metastatic colorectal cancer (mCRC) patients enrolled in FIRE-3 study. *Journal of Clinical Oncology* 34, 3516-3516.
- Le, D.T., Durham, J.N., Smith, K.N., Wang, H., Bartlett, B.R., Aulakh, L.K., Lu, S., Kemberling, H., Wilt, C., Lubner, B.S., Wong, F., Azad, N.S., Rucki, A.A., Laheru, D., Donehower, R., Zaheer, A., Fisher, G.A., Crocenzi, T.S., Lee, J.J., Greten, T.F., Duffy, A.G., Ciombor, K.K., Eyring, A.D., Lam, B.H., Joe, A., Kang, S.P., Holdhoff, M., Danilova, L., Cope, L., Meyer, C., Zhou, S., Goldberg, R.M., Armstrong, D.K., Bever, K.M., Fader, A.N., Taube, J., Housseau, F., Spetzler, D., Xiao, N., Pardoll, D.M., Papadopoulos, N., Kinzler, K.W., Eshleman, J.R., Vogelstein, B., Anders, R.A., and Diaz, L.A., Jr. (2017). Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 357, 409-413.
- Le Gall, M., Chambard, J.C., Breitmayer, J.P., Grall, D., Pouyssegur, J., and Van Obberghen-Schilling, E. (2000). The p42/p44 MAP kinase pathway prevents apoptosis induced by anchorage and serum removal. *Mol Biol Cell* 11, 1103-1112.
- Lee, C.G., McCarthy, S., Gruidl, M., Timme, C., and Yeatman, T.J. (2014). MicroRNA-147 Induces a Mesenchymal-To-Epithelial Transition (MET) and Reverses EGFR Inhibitor Resistance. *Plos One* 9.
- Li, B.S., Zuo, Q.F., Zhao, Y.L., Xiao, B., Zhuang, Y., Mao, X.H., Wu, C., Yang, S.M., Zeng, H., Zou, Q.M., and Guo, G. (2015a). MicroRNA-25 promotes gastric cancer migration, invasion and proliferation by directly targeting transducer of ERBB2, 1 and correlates with poor survival. *Oncogene* 34, 2556-2565.
- Li, C., Liang, G., Yang, S., Sui, J., Yao, W., Shen, X., Zhang, Y., Peng, H., Hong, W., Xu, S., Wu, W., Ye, Y., Zhang, Z., Zhang, W., Yin, L., and Pu, Y. (2017). Dysregulated lncRNA-UCA1 contributes to the progression of gastric cancer through regulation of the PI3K-Akt-mTOR signaling pathway. *Oncotarget* 8, 93476-93491.
- Li, C.H., and Chen, Y. (2016). Insight Into the Role of Long Noncoding RNA in Cancer Development and Progression. *Int Rev Cell Mol Biol* 326, 33-65.
- Li, F., Mao, G., Tong, D., Huang, J., Gu, L., Yang, W., and Li, G.M. (2013). The histone mark H3K36me3 regulates human DNA mismatch repair through its interaction with MutSalpha. *Cell* 153, 590-600.
- Li, G., Liu, C., Yuan, J., Xiao, X., Tang, N., Hao, J., Wang, H., Bian, X., Deng, Y., and Ding, Y. (2010). CD133(+) single cell-derived progenies of colorectal cancer cell line SW480 with different invasive and metastatic potential. *Clin Exp Metastasis* 27, 517-527.
- Li, G., Zhang, H., Wan, X., Yang, X., Zhu, C., Wang, A., He, L., Miao, R., Chen, S., and Zhao, H. (2014a). Long noncoding RNA plays a key role in metastasis and prognosis of hepatocellular carcinoma. *Biomed Res Int* 2014, 780521.
- Li, H., An, J., Wu, M., Zheng, Q., Gui, X., Li, T., Pu, H., and Lu, D. (2015b). LncRNA HOTAIR promotes human liver cancer stem cell malignant growth through downregulation of SETD2. *Oncotarget* 6, 27847-27864.

- Li, J., Zhang, Y., Zhao, J., Kong, F., and Chen, Y. (2011). Overexpression of miR-22 reverses paclitaxel-induced chemoresistance through activation of PTEN signaling in p53-mutated colon cancer cells. *Mol Cell Biochem* 357, 31-38.
- Li, J.Y., Ma, X., and Zhang, C.B. (2014b). Overexpression of long non-coding RNA UCA1 predicts a poor prognosis in patients with esophageal squamous cell carcinoma. *Int J Clin Exp Pathol* 7, 7938-7944.
- Li, T., Zheng, Q., An, J., Wu, M., Li, H., Gui, X., Pu, H., and Lu, D. (2016a). SET1A Cooperates With CUDR to Promote Liver Cancer Growth and Hepatocyte-like Stem Cell Malignant Transformation Epigenetically. *Mol Ther* 24, 261-275.
- Li, X., Li, X., Liao, D., Wang, X., Wu, Z., Nie, J., Bai, M., Fu, X., Mei, Q., and Han, W. (2015c). Elevated microRNA-23a Expression Enhances the Chemoresistance of Colorectal Cancer Cells with Microsatellite Instability to 5-Fluorouracil by Directly Targeting ABCF1. *Curr Protein Pept Sci* 16, 301-309.
- Li, X., Li, X.L., Liao, D.X., Wang, X.H., Wu, Z.Q., Nie, J., Bai, M.M., Fu, X.B., Mei, Q., and Han, W.D. (2015d). Elevated microRNA-23a Expression Enhances the Chemoresistance of Colorectal Cancer Cells with Microsatellite Instability to 5-Fluorouracil by Directly Targeting ABCF1. *Current Protein & Peptide Science* 16, 301-309.
- Li, Y., Vandenboom, T.G., 2nd, Kong, D., Wang, Z., Ali, S., Philip, P.A., and Sarkar, F.H. (2009). Up-regulation of miR-200 and let-7 by natural agents leads to the reversal of epithelial-to-mesenchymal transition in gemcitabine-resistant pancreatic cancer cells. *Cancer Res* 69, 6704-6712.
- Li, Z., Shen, J.X., Chan, M.T.V., and Wu, W.K.K. (2016b). TUG1: a pivotal oncogenic long non-coding RNA of human cancers. *Cell Proliferation* 49, 471-475.
- Li, Z., Zhao, X., Zhou, Y., Liu, Y., Zhou, Q., Ye, H., Wang, Y., Zeng, J., Song, Y., Gao, W., Zheng, S., Zhuang, B., Chen, H., Li, W., Li, H., Li, H., Fu, Z., and Chen, R. (2015e). The long non-coding RNA HOTTIP promotes progression and gemcitabine resistance by regulating HOXA13 in pancreatic cancer. *J Transl Med* 13, 84.
- Lim, L.P., Lau, N.C., Garrett-Engele, P., Grimson, A., Schelter, J.M., Castle, J., Bartel, D.P., Linsley, P.S., and Johnson, J.M. (2005). Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 433, 769-773.
- Lin, C.P., Liu, J.D., Chow, J.M., Liu, C.R., and Liu, H.E. (2007). Small-molecule c-Myc inhibitor, 10058-F4, inhibits proliferation, downregulates human telomerase reverse transcriptase and enhances chemosensitivity in human hepatocellular carcinoma cells. *Anticancer Drugs* 18, 161-170.
- Lin, C.Y., and Xu, H.M. (2015). Novel perspectives of long non-coding RNAs in esophageal carcinoma. *Carcinogenesis* 36, 1255-1262.
- Lin, H.M., Mahon, K.L., Spielman, C., Gurney, H., Mallesara, G., Stockler, M.R., Bastick, P., Briscoe, K., Marx, G., Swarbrick, A., and Horvath, L.G. (2017). Phase 2 study of circulating microRNA biomarkers in castration-resistant prostate cancer. *Br J Cancer* 116, 1002-1011.
- Lin, L., Yee, S.W., Kim, R.B., and Giacomini, K.M. (2015). SLC transporters as therapeutic targets: emerging opportunities. *Nature Reviews Drug Discovery* 14, 543-560.
- Lin, M., Gu, J., Eng, C., Ellis, L.M., Hildebrandt, M.A., Lin, J., Huang, M., Calin, G.A., Wang, D., Dubois, R.N., Hawk, E.T., and Wu, X. (2012). Genetic

- polymorphisms in MicroRNA-related genes as predictors of clinical outcomes in colorectal adenocarcinoma patients. *Clin Cancer Res* 18, 3982-3991.
- Ling, H., Fabbri, M., and Calin, G.A. (2013). MicroRNAs and other non-coding RNAs as targets for anticancer drug development. *Nat Rev Drug Discov* 12, 847-865.
- Liu, C., and Tang, D.G. (2011). MicroRNA regulation of cancer stem cells. *Cancer Res* 71, 5950-5954.
- Liu, Z.L., Sun, M., Lu, K.H., Liu, J., Zhang, M.L., Wu, W.Q., De, W., Wang, Z.X., and Wang, R. (2013). The Long Noncoding RNA HOTAIR Contributes to Cisplatin Resistance of Human Lung Adenocarcinoma Cells via downregulation of p21(WAF1/CIP1) Expression. *Plos One* 8.
- Longley, D.B., and Johnston, P.G. (2005). Molecular mechanisms of drug resistance. *J Pathol* 205, 275-292.
- Lorusso, P.M. (2016). Inhibition of the PI3K/AKT/mTOR Pathway in Solid Tumors. *J Clin Oncol* 34, 3803-3815.
- Loupakis, F., Bria, E., Vaccaro, V., Cuppone, F., Milella, M., Carlini, P., Cremolini, C., Salvatore, L., Falcone, A., Muti, P., Sperduti, I., Giannarelli, D., and Cognetti, F. (2010). Magnitude of benefit of the addition of bevacizumab to first-line chemotherapy for metastatic colorectal cancer: meta-analysis of randomized clinical trials. *J Exp Clin Cancer Res* 29, 58.
- Lu, M., Liu, Z., Li, B., Wang, G., Li, D.C., and Zhu, Y.P. (2017a). The high expression of long non-coding RNA PANDAR indicates a poor prognosis for colorectal cancer and promotes metastasis by EMT pathway. *Journal of Cancer Research and Clinical Oncology* 143, 71-81.
- Lu, Y., Zhao, X., Liu, Q., Li, C., Graves-Deal, R., Cao, Z., Singh, B., Franklin, J.L., Wang, J., Hu, H., Wei, T., Yang, M., Yeatman, T.J., Lee, E., Saito-Diaz, K., Hinger, S., Patton, J.G., Chung, C.H., Emmrich, S., Klusmann, J.H., Fan, D., and Coffey, R.J. (2017b). lncRNA MIR100HG-derived miR-100 and miR-125b mediate cetuximab resistance via Wnt/beta-catenin signaling. *Nat Med* 23, 1331-1341.
- Ma, J., Fang, B., Zeng, F., Ma, C., Pang, H., Cheng, L., Shi, Y., Wang, H., Yin, B., Xia, J., and Wang, Z. (2015). Down-regulation of miR-223 reverses epithelial-mesenchymal transition in gemcitabine-resistant pancreatic cancer cells. *Oncotarget* 6, 1740-1749.
- Ma, P., Xu, T.P., Huang, M.D., and Shu, Y.Q. (2016a). Increased expression of LncRNA PANDAR predicts a poor prognosis in gastric cancer. *Biomedicine & Pharmacotherapy* 78, 172-176.
- Ma, S., Tang, K.H., Chan, Y.P., Lee, T.K., Kwan, P.S., Castilho, A., Ng, I., Man, K., Wong, N., To, K.F., Zheng, B.J., Lai, P.B., Lo, C.M., Chan, K.W., and Guan, X.Y. (2010). miR-130b Promotes CD133(+) liver tumor-initiating cell growth and self-renewal via tumor protein 53-induced nuclear protein 1. *Cell Stem Cell* 7, 694-707.
- Ma, Y.L., Yang, Y.Z., Wang, F., Moyer, M.P., Wei, Q., Zhang, P., Yang, Z., Liu, W.J., Zhang, H.Z., Chen, N.W., Wang, H., Wang, H.M., and Qin, H.L. (2016b). Long non-coding RNA CCAL regulates colorectal cancer progression by activating Wnt/beta-catenin signalling pathway via suppression of activator protein 2 alpha. *Gut* 65, 1494-1504.

- Macfarlane, L.A., and Murphy, P.R. (2010). MicroRNA: Biogenesis, Function and Role in Cancer. *Curr Genomics* 11, 537-561.
- Maemondo, M., Inoue, A., Kobayashi, K., Sugawara, S., Oizumi, S., Isobe, H., Gemma, A., Harada, M., Yoshizawa, H., Kinoshita, I., Fujita, Y., Okinaga, S., Hirano, H., Yoshimori, K., Harada, T., Ogura, T., Ando, M., Miyazawa, H., Tanaka, T., Saijo, Y., Hagiwara, K., Morita, S., Nukiwa, T., and North-East Japan Study, G. (2010). Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med* 362, 2380-2388.
- Maftouh, M., Avan, A., Funel, N., Frampton, A.E., Fiuji, H., Pelliccioni, S., Castellano, L., Galla, V., Peters, G.J., and Giovannetti, E. (2014). Mir-211 Modulates Gemcitabine Activity through Downregulation of Ribonucleotide Reductase and Inhibits the Invasive Behavior of Pancreatic Cancer Cells. *Nucleosides Nucleotides & Nucleic Acids* 33, 384-393.
- Matouk, I., Raveh, E., Ohana, P., Abu Lail, R., Gershtain, E., Gilon, M., De Groot, N., Czerniak, A., and Hochberg, A. (2013). The Increasing Complexity of the Oncofetal H19 Gene Locus: Functional Dissection and Therapeutic Intervention. *International Journal of Molecular Sciences* 14, 4298-4316.
- Mckenna, L.B., Schug, J., Vourekas, A., Mckenna, J.B., Bramswig, N.C., Friedman, J.R., and Kaestner, K.H. (2010). MicroRNAs Control Intestinal Epithelial Differentiation, Architecture, and Barrier Function. *Gastroenterology* 139, 1654-+.
- Mclellan, L.I., and Wolf, C.R. (1999a). Glutathione and glutathione-dependent enzymes in cancer drug resistance. *Drug Resistance Updates* 2, 153-164.
- Mclellan, L.I., and Wolf, C.R. (1999b). Glutathione and glutathione-dependent enzymes in cancer drug resistance. *Drug Resist Updat* 2, 153-164.
- Mebratu, Y., and Tesfagzi, Y. (2009). How ERK1/2 activation controls cell proliferation and cell death: Is subcellular localization the answer? *Cell Cycle* 8, 1168-1175.
- Meng, F., Henson, R., Wehbe-Jane, H., Ghoshal, K., Jacob, S.T., and Patel, T. (2007). MicroRNA-21 regulates expression of the PTEN tumor suppressor gene in human hepatocellular cancer. *Gastroenterology* 133, 647-658.
- Michael, M., and Doherty, M.M. (2005). Tumoral drug metabolism: overview and its implications for cancer therapy. *J Clin Oncol* 23, 205-229.
- Mitchell, P.S., Parkin, R.K., Kroh, E.M., Fritz, B.R., Wyman, S.K., Pogosova-Agadjanyan, E.L., Peterson, A., Noteboom, J., O'briant, K.C., Allen, A., Lin, D.W., Urban, N., Drescher, C.W., Knudsen, B.S., Stirewalt, D.L., Gentleman, R., Vessella, R.L., Nelson, P.S., Martin, D.B., and Tewari, M. (2008). Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A* 105, 10513-10518.
- Mitra, A., Mishra, L., and Li, S. (2015). EMT, CTCs and CSCs in tumor relapse and drug-resistance. *Oncotarget* 6, 10697-10711.
- Mohamadkhani, A. (2014). Long Noncoding RNAs in Interaction With RNA Binding Proteins in Hepatocellular Carcinoma. *Hepat Mon* 14, e18794.
- Mohri, T., Nakajima, M., Fukami, T., Takamiya, M., Aoki, Y., and Yokoi, T. (2010). Human CYP2E1 is regulated by miR-378. *Biochemical Pharmacology* 79, 1045-1052.

- Moran, V.A., Perera, R.J., and Khalil, A.M. (2012). Emerging functional and mechanistic paradigms of mammalian long non-coding RNAs. *Nucleic Acids Res* 40, 6391-6400.
- Moriyama, T., Ohuchida, K., Mizumoto, K., Yu, J., Sato, N., Nabae, T., Takahata, S., Toma, H., Nagai, E., and Tanaka, M. (2009). MicroRNA-21 modulates biological functions of pancreatic cancer cells including their proliferation, invasion, and chemoresistance. *Mol Cancer Ther* 8, 1067-1074.
- Mossink, M.H., Van Zon, A., Scheper, R.J., Sonneveld, P., and Wiemer, E.a.C. (2003). Vaults: a ribonucleoprotein particle involved in drug resistance? *Oncogene* 22, 7458-7467.
- Motzer, R.J., Hutson, T.E., Tomczak, P., Michaelson, M.D., Bukowski, R.M., Rixe, O., Oudard, S., Negrier, S., Szczylik, C., Kim, S.T., Chen, I., Bycott, P.W., Baum, C.M., and Figlin, R.A. (2007). Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med* 356, 115-124.
- Munoz, M., Henderson, M., Haber, M., and Norris, M. (2007). Role of the MRP1/ABCC1 multidrug transporter protein in cancer. *Lab Invest* 87, 752-757.
- Murakami, Y., Yasuda, T., Saigo, K., Urashima, T., Toyoda, H., Okanoue, T., and Shimotohno, K. (2006). Comprehensive analysis of microRNA expression patterns in hepatocellular carcinoma and non-tumorous tissues. *Oncogene* 25, 2537-2545.
- Muro, K., Bang, Y.J., Shankaran, V., Geva, R., Catenacci, D.V.T., Gupta, S., Eder, J.P., Berger, R., Gonzalez, E.J., Ray, A., Dolled-Filhart, M., Emancipator, K., Pathiraja, K., Lunceford, J.K., Cheng, J.D., Koshiji, M., and Chung, H.C. (2015). Relationship between PD-L1 expression and clinical outcomes in patients (Pts) with advanced gastric cancer treated with the anti-PD-1 monoclonal antibody pembrolizumab (Pembro; MK-3475) in KEYNOTE-012. *Journal of Clinical Oncology* 33.
- Murphy, M.B. (2016). Adjunctive Therapy of Gastric Cancer: Moving the Field Forward. *Oncology (Williston Park)* 30, 646-647.
- Nguyen, L.V., Vanner, R., Dirks, P., and Eaves, C.J. (2012). Cancer stem cells: an evolving concept. *Nat Rev Cancer* 12, 133-143.
- Nho, R.S., and Hergert, P. (2014). FoxO3a and disease progression. *World J Biol Chem* 5, 346-354.
- Nie, J., Liu, L., Zheng, W., Chen, L., Wu, X., Xu, Y., Du, X., and Han, W. (2012). microRNA-365, down-regulated in colon cancer, inhibits cell cycle progression and promotes apoptosis of colon cancer cells by probably targeting Cyclin D1 and Bcl-2. *Carcinogenesis* 33, 220-225.
- Niinuma, T., Suzuki, H., Nojima, M., Noshio, K., Yamamoto, H., Takamaru, H., Yamamoto, E., Maruyama, R., Nobuoka, T., Miyazaki, Y., Nishida, T., Bamba, T., Kanda, T., Ajioka, Y., Taguchi, T., Okahara, S., Takahashi, H., Nishida, Y., Hosokawa, M., Hasegawa, T., Tokino, T., Hirata, K., Imai, K., Toyota, M., and Shinomura, Y. (2012). Upregulation of miR-196a and HOTAIR Drive Malignant Character in Gastrointestinal Stromal Tumors. *Cancer Research* 72, 1126-1136.
- Nishida, N., Yamashita, S., Mimori, K., Sudo, T., Tanaka, F., Shibata, K., Yamamoto, H., Ishii, H., Doki, Y., and Mori, M. (2012). MicroRNA-10b is a prognostic

- indicator in colorectal cancer and confers resistance to the chemotherapeutic agent 5-fluorouracil in colorectal cancer cells. *Ann Surg Oncol* 19, 3065-3071.
- Nurwidya, F., Takahashi, F., Murakami, A., and Takahashi, K. (2012). Epithelial mesenchymal transition in drug resistance and metastasis of lung cancer. *Cancer Res Treat* 44, 151-156.
- O'connell, R.M., Rao, D.S., Chaudhuri, A.A., and Baltimore, D. (2010). Physiological and pathological roles for microRNAs in the immune system. *Nat Rev Immunol* 10, 111-122.
- Offer, S.M., Butterfield, G.L., Jerde, C.R., Fossum, C.C., Wegner, N.J., and Diasio, R.B. (2014). microRNAs miR-27a and miR-27b Directly Regulate Liver Dihydropyrimidine Dehydrogenase Expression through Two Conserved Binding Sites. *Molecular Cancer Therapeutics* 13, 742-751.
- Olcina, M.M., and Giaccia, A.J. (2016). Reducing radiation-induced gastrointestinal toxicity - the role of the PHD/HIF axis. *J Clin Invest* 126, 3708-3715.
- Ounzain, S., Pezzuto, I., Micheletti, R., Burdet, F., Sheta, R., Nemir, M., Gonzales, C., Sarre, A., Alexanian, M., Blow, M.J., May, D., Johnson, R., Dauvillier, J., Pennacchio, L.A., and Pedrazzini, T. (2014). Functional importance of cardiac enhancer-associated noncoding RNAs in heart development and disease. *J Mol Cell Cardiol* 76, 55-70.
- Overman, M.J., Lonardi, S., Wong, K.Y.M., Lenz, H.J., Gelsomino, F., Aglietta, M., Morse, M.A., Van Cutsem, E., Mcdermott, R., Hill, A., Sawyer, M.B., Hendlish, A., Neyns, B., Svrcek, M., Moss, R.A., Ledine, J.M., Cao, Z.A., Kamble, S., Kopetz, S., and Andre, T. (2018). Durable Clinical Benefit With Nivolumab Plus Ipilimumab in DNA Mismatch Repair-Deficient/Microsatellite Instability-High Metastatic Colorectal Cancer. *J Clin Oncol* 36, 773-779.
- Ozawa, T., Kandimalla, R., Gao, F., Nozawa, H., Hata, K., Nagata, H., Okada, S., Izumi, D., Baba, H., Fleshman, J., Wang, X., Watanabe, T., and Goel, A. (2018). A MicroRNA Signature Associated With Metastasis of T1 Colorectal Cancers to Lymph Nodes. *Gastroenterology* 154, 844-848 e847.
- Padua Alves, C., Fonseca, A.S., Muys, B.R., De Barros, E.L.B.R., Burger, M.C., De Souza, J.E., Valente, V., Zago, M.A., and Silva, W.A., Jr. (2013). Brief report: The lincRNA Hotair is required for epithelial-to-mesenchymal transition and stemness maintenance of cancer cell lines. *Stem Cells* 31, 2827-2832.
- Pan, Y.Z., Gao, W.Q., and Yu, A.M. (2009). MicroRNAs Regulate CYP3A4 Expression via Direct and Indirect Targeting. *Drug Metabolism and Disposition* 37, 2112-2117.
- Pan, Y.Z., Zhou, A., Hu, Z.H., and Yu, A.M. (2013). Small Nucleolar RNA-Derived MicroRNA hsa-miR-1291 Modulates Cellular Drug Disposition through Direct Targeting of ABC Transporter ABCC1. *Drug Metabolism and Disposition* 41, 1744-1751.
- Pardini, B., Rosa, F., Naccarati, A., Vymetalkova, V., Ye, Y., Wu, X., Di Gaetano, C., Buchler, T., Novotny, J., Matullo, G., and Vodicka, P. (2015). Polymorphisms in microRNA genes as predictors of clinical outcomes in colorectal cancer patients. *Carcinogenesis* 36, 82-86.
- Park, G.R. (2001). Sedation and analgesia-which way is best? *Br J Anaesth* 87, 183-185.

- 1699 Park, J.K., Lee, E.J., Esau, C., and Schmittgen, T.D. (2009). Antisense inhibition of
1700 microRNA-21 or -221 arrests cell cycle, induces apoptosis, and sensitizes the
1701 effects of gemcitabine in pancreatic adenocarcinoma. *Pancreas* 38, e190-199.
- 1702 Pasquinelli, A.E. (2012). MicroRNAs and their targets: recognition, regulation and an
1703 emerging reciprocal relationship. *Nat Rev Genet* 13, 271-282.
- 1704 Pattabiraman, D.R., and Weinberg, R.A. (2014). Tackling the cancer stem cells -
1705 what challenges do they pose? *Nat Rev Drug Discov* 13, 497-512.
- 1706 Pearl, L.H., Schierz, A.C., Ward, S.E., Al-Lazikani, B., and Pearl, F.M.G. (2015).
1707 Therapeutic opportunities within the DNA damage response. *Nature Reviews*
1708 *Cancer* 15, 166-180.
- 1709 Peng, W., and Fan, H. (2015). Long non-coding RNA PANDAR correlates with poor
1710 prognosis and promotes tumorigenesis in hepatocellular carcinoma.
1711 *Biomedicine & Pharmacotherapy* 72, 113-118.
- 1712 Persson, H., Kvist, A., Vallon-Christersson, J., Medstrand, P., Borg, A., and Rovira,
1713 C. (2009). The non-coding RNA of the multidrug resistance-linked vault
1714 particle encodes multiple regulatory small RNAs. *Nature Cell Biology* 11,
1715 1268-U1265.
- 1716 Peterson, L.W., and Artis, D. (2014). Intestinal epithelial cells: regulators of barrier
1717 function and immune homeostasis. *Nature Reviews Immunology* 14, 141-153.
- 1718 Pfister, S.X., Ahrabi, S., Zalmas, L.P., Sarkar, S., Aymard, F., Bachrati, C.Z.,
1719 Helleday, T., Legube, G., La Thangue, N.B., Porter, A.C., and Humphrey, T.C.
1720 (2014). SETD2-dependent histone H3K36 trimethylation is required for
1721 homologous recombination repair and genome stability. *Cell Rep* 7, 2006-
1722 2018.
- 1723 Pothof, J., Verkaik, N.S., Van Ijcken, W., Wiemer, E.a.C., Ta, V.T.B., Van Der Horst,
1724 G.T.J., Jaspers, N.G.J., Van Gent, D.C., Hoeijmakers, J.H.J., and Persengiev,
1725 S.P. (2009). MicroRNA-mediated gene silencing modulates the UV-induced
1726 DNA-damage response. *Embo Journal* 28, 2090-2099.
- 1727 Pourhoseingholi, M.A., Vahedi, M., and Baghestani, A.R. (2015). Burden of
1728 gastrointestinal cancer in Asia; an overview. *Gastroenterol Hepatol Bed*
1729 *Bench* 8, 19-27.
- 1730 Prieto-Vila, M., Takahashi, R.U., Usuba, W., Kohama, I., and Ochiya, T. (2017). Drug
1731 Resistance Driven by Cancer Stem Cells and Their Niche. *Int J Mol Sci* 18.
- 1732 Pu, H., Zheng, Q., Li, H., Wu, M., An, J., Gui, X., Li, T., and Lu, D. (2015). CUDR
1733 promotes liver cancer stem cell growth through upregulating TERT and C-
1734 Myc. *Oncotarget* 6, 40775-40798.
- 1735 Pullen, T.J., Xavier, G.D., Kelsey, G., and Rutter, G.A. (2011). miR-29a and miR-29b
1736 Contribute to Pancreatic beta-Cell-Specific Silencing of Monocarboxylate
1737 Transporter 1 (Mct1). *Molecular and Cellular Biology* 31, 3182-3194.
- 1738 Pyndiah, S., Tanida, S., Ahmed, K.M., Cassimere, E.K., Choe, C., and Sakamuro, D.
1739 (2011). c-MYC suppresses BIN1 to release poly(ADP-ribose) polymerase 1: a
1740 mechanism by which cancer cells acquire cisplatin resistance. *Sci Signal* 4,
1741 ra19.
- 1742 Qi, P., Zhou, X.Y., and Du, X. (2016). Circulating long non-coding RNAs in cancer:
1743 current status and future perspectives. *Mol Cancer* 15, 39.
- 1744 Qu, J., Zhao, L., Zhang, P., Wang, J., Xu, N., Mi, W., Jiang, X., Zhang, C., and Qu, J.
1745 (2015). MicroRNA-195 chemosensitizes colon cancer cells to the

- chemotherapeutic drug doxorubicin by targeting the first binding site of BCL2L2 mRNA. *J Cell Physiol* 230, 535-545.
- Quinn, J.J., and Chang, H.Y. (2016). Unique features of long non-coding RNA biogenesis and function. *Nat Rev Genet* 17, 47-62.
- Rautio, T., Kairaluoma, M., and Sand, J. (2016). [Novel techniques in the treatment of rectal cancer]. *Duodecim* 132, 1160-1164.
- Rendic, S., and Guengerich, F.P. (2015). Survey of Human Oxidoreductases and Cytochrome P450 Enzymes Involved in the Metabolism of Xenobiotic and Natural Chemicals. *Chem Res Toxicol* 28, 38-42.
- Reya, T., Morrison, S.J., Clarke, M.F., and Weissman, I.L. (2001). Stem cells, cancer, and cancer stem cells. *Nature* 414, 105-111.
- Ristamaki, R., and Algars, A. (2016). [Principles of oncologic drug therapy following surgery for bowel cancer]. *Duodecim* 132, 1155-1159.
- Rodrigues, A.S., Dinis, J., Gromicho, M., Martins, C., Laires, A., and Rueff, J. (2012a). Genomics and cancer drug resistance. *Curr Pharm Biotechnol* 13, 651-673.
- Rodrigues, A.S., Dinis, J., Gromicho, M., Martins, C., Laires, A., and Rueff, J. (2012b). Genomics and Cancer Drug Resistance. *Current Pharmaceutical Biotechnology* 13, 651-673.
- Roukos, D.H. (2010). Targeting gastric cancer with trastuzumab: new clinical practice and innovative developments to overcome resistance. *Ann Surg Oncol* 17, 14-17.
- Rupaimoole, R., and Slack, F.J. (2017). MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat Rev Drug Discov* 16, 203-222.
- Rutkowski, P., and Hompes, D. (2016). Combined Therapy of Gastrointestinal Stromal Tumors. *Surg Oncol Clin N Am* 25, 735-759.
- Sacconi, A., Biagioni, F., Canu, V., Mori, F., Di Benedetto, A., Lorenzon, L., Ercolani, C., Di Agostino, S., Cambria, A.M., Germoni, S., Grasso, G., Blandino, R., Panebianco, V., Ziparo, V., Federici, O., Muti, P., Strano, S., Carboni, F., Mottolese, M., Diodoro, M., Pescarmona, E., Garofalo, A., and Blandino, G. (2012). miR-204 targets Bcl-2 expression and enhances responsiveness of gastric cancer. *Cell Death Dis* 3, e423.
- Safa, A.R. (2016). Resistance to Cell Death and Its Modulation in Cancer Stem Cells. *Crit Rev Oncog* 21, 203-219.
- Sasaki, M., Ikeda, H., Itatsu, K., Yamaguchi, J., Sawada, S., Minato, H., Ohta, T., and Nakanuma, Y. (2008). The overexpression of polycomb group proteins Bmi1 and EZH2 is associated with the progression and aggressive biological behavior of hepatocellular carcinoma. *Lab Invest* 88, 873-882.
- Sauvageau, M., Goff, L.A., Lodato, S., Bonev, B., Groff, A.F., Gerhardinger, C., Sanchez-Gomez, D.B., Hacisuleyman, E., Li, E., Spence, M., Liapis, S.C., Mallard, W., Morse, M., Swerdel, M.R., D'ecclissis, M.F., Moore, J.C., Lai, V., Gong, G., Yancopoulos, G.D., Frendewey, D., Kellis, M., Hart, R.P., Valenzuela, D.M., Arlotta, P., and Rinn, J.L. (2013). Multiple knockout mouse models reveal lincRNAs are required for life and brain development. *Elife* 2, e01749.

- Schreiber, R., Mezencev, R., Matyunina, L.V., and McDonald, J.F. (2016). Evidence for the role of microRNA 374b in acquired cisplatin resistance in pancreatic cancer cells. *Cancer Gene Therapy* 23, 241-245.
- Schultz, N.A., Dehlendorff, C., Jensen, B.V., Bjerregaard, J.K., Nielsen, K.R., Bojesen, S.E., Calatayud, D., Nielsen, S.E., Yilmaz, M., Hollander, N.H., Andersen, K.K., and Johansen, J.S. (2014). MicroRNA biomarkers in whole blood for detection of pancreatic cancer. *JAMA* 311, 392-404.
- Sclafani, F., Chau, I., Cunningham, D., Lampis, A., Hahne, J.C., Ghidini, M., Lote, H., Zito, D., Tabernero, J., Glimelius, B., Cervantes, A., Begum, R., De Castro, D.G., Wilson, S.H., Peckitt, C., Eltahir, Z., Wotherspoon, A., Tait, D., Brown, G., Oates, J., Braconi, C., and Valeri, N. (2016). Sequence variation in mature microRNA-608 and benefit from neo-adjuvant treatment in locally advanced rectal cancer patients. *Carcinogenesis* 37, 852-857.
- Sclafani, F., Chau, I., Cunningham, D., Peckitt, C., Lampis, A., Hahne, J.C., Braconi, C., Tabernero, J., Glimelius, B., Cervantes, A., Begum, R., Gonzalez De Castro, D., Hulkki Wilson, S., Eltahir, Z., Wotherspoon, A., Tait, D., Brown, G., Oates, J., and Valeri, N. (2015). Prognostic role of the LCS6 KRAS variant in locally advanced rectal cancer: results of the EXPERT-C trial. *Ann Oncol* 26, 1936-1941.
- Sha, D., Lee, A.M., Shi, Q., Alberts, S.R., Sargent, D.J., Sinicrope, F.A., and Diasio, R.B. (2014). Association study of the let-7 miRNA-complementary site variant in the 3' untranslated region of the KRAS gene in stage III colon cancer (NCCTG N0147 Clinical Trial). *Clin Cancer Res* 20, 3319-3327.
- Shang, C., Guo, Y., Zhang, J., and Huang, B. (2016). Silence of long noncoding RNA UCA1 inhibits malignant proliferation and chemotherapy resistance to adriamycin in gastric cancer. *Cancer Chemother Pharmacol* 77, 1061-1067.
- Shang, Y., Cai, X., and Fan, D. (2013). Roles of epithelial-mesenchymal transition in cancer drug resistance. *Curr Cancer Drug Targets* 13, 915-929.
- Shang, Y., Zhang, Z., Liu, Z., Feng, B., Ren, G., Li, K., Zhou, L., Sun, Y., Li, M., Zhou, J., An, Y., Wu, K., Nie, Y., and Fan, D. (2014). miR-508-5p regulates multidrug resistance of gastric cancer by targeting ABCB1 and ZNRD1. *Oncogene* 33, 3267-3276.
- Shankar, S., Nall, D., Tang, S.N., Meeker, D., Passarini, J., Sharma, J., and Srivastava, R.K. (2011). Resveratrol inhibits pancreatic cancer stem cell characteristics in human and KrasG12D transgenic mice by inhibiting pluripotency maintaining factors and epithelial-mesenchymal transition. *PLoS One* 6, e16530.
- Sharma, V., and Misteli, T. (2013). Non-coding RNAs in DNA damage and repair. *Febs Letters* 587, 1832-1839.
- Shea, T.C., Kelley, S.L., and Henner, W.D. (1988a). Identification of an Anionic Form of Glutathione Transferase Present in Many Human-Tumors and Human-Tumor Cell-Lines. *Cancer Research* 48, 527-533.
- Shea, T.C., Kelley, S.L., and Henner, W.D. (1988b). Identification of an anionic form of glutathione transferase present in many human tumors and human tumor cell lines. *Cancer Res* 48, 527-533.
- Shigeyasu, K., Toden, S., Zumwalt, T.J., Okugawa, Y., and Goel, A. (2017). Emerging Role of MicroRNAs as Liquid Biopsy Biomarkers in Gastrointestinal Cancers. *Clin Cancer Res* 23, 2391-2399.

- Shimada, T., Iwasaki, M., Martin, M.V., and Guengerich, F.P. (1989). Human liver microsomal cytochrome P-450 enzymes involved in the bioactivation of procarcinogens detected by umu gene response in *Salmonella typhimurium* TA 1535/pSK1002. *Cancer Res* 49, 3218-3228.
- Siegel, R.L., Miller, K.D., and Jemal, A. (2015). Cancer statistics, 2015. *CA Cancer J Clin* 65, 5-29.
- Singh, S., Chitkara, D., Kumar, V., Behrman, S.W., and Mahato, R.I. (2013). miRNA profiling in pancreatic cancer and restoration of chemosensitivity. *Cancer Lett* 334, 211-220.
- Siroy, A.E., Boland, G.M., Milton, D.R., Roszik, J., Frankian, S., Malke, J., Haydu, L., Prieto, V.G., Tetzlaff, M., Ivan, D., Wang, W.L., Torres-Cabala, C., Curry, J., Roy-Chowdhuri, S., Broaddus, R., Rashid, A., Stewart, J., Gershenwald, J.E., Amaria, R.N., Patel, S.P., Papadopoulos, N.E., Bedikian, A., Hwu, W.J., Hwu, P., Diab, A., Woodman, S.E., Aldape, K.D., Luthra, R., Patel, K.P., Shaw, K.R., Mills, G.B., Mendelsohn, J., Meric-Bernstam, F., Kim, K.B., Routbort, M.J., Lazar, A.J., and Davies, M.A. (2015). Beyond BRAF(V600): clinical mutation panel testing by next-generation sequencing in advanced melanoma. *J Invest Dermatol* 135, 508-515.
- Slamon, D.J., Leyland-Jones, B., Shak, S., Fuchs, H., Paton, V., Bajamonde, A., Fleming, T., Eiermann, W., Wolter, J., Pegram, M., Baselga, J., and Norton, L. (2001). Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 344, 783-792.
- Smith, J., Tho, L.M., Xu, N., and Gillespie, D.A. (2010). The ATM-Chk2 and ATR-Chk1 pathways in DNA damage signaling and cancer. *Adv Cancer Res* 108, 73-112.
- Smolle, M.A., Calin, H.N., Pichler, M., and Calin, G.A. (2017). Noncoding RNAs and immune checkpoints-clinical implications as cancer therapeutics. *FEBS J* 284, 1952-1966.
- Song, J., Qu, Z., Guo, X., Zhao, Q., Zhao, X., Gao, L., Sun, K., Shen, F., Wu, M., and Wei, L. (2009). Hypoxia-induced autophagy contributes to the chemoresistance of hepatocellular carcinoma cells. *Autophagy* 5, 1131-1144.
- Srivastava, R.K., Tang, S.N., Zhu, W., Meeker, D., and Shankar, S. (2011). Sulforaphane synergizes with quercetin to inhibit self-renewal capacity of pancreatic cancer stem cells. *Front Biosci (Elite Ed)* 3, 515-528.
- Sui, M., Jiao, A., Zhai, H., Wang, Y., Wang, Y., Sun, D., and Li, P. (2017). Upregulation of miR-125b is associated with poor prognosis and trastuzumab resistance in HER2-positive gastric cancer. *Exp Ther Med* 14, 657-663.
- Sun, X., Jiao, X., Pestell, T.G., Fan, C., Qin, S., Mirabelli, E., Ren, H., and Pestell, R.G. (2014). MicroRNAs and cancer stem cells: the sword and the shield. *Oncogene* 33, 4967-4977.
- Taal, B.G., Audisio, R.A., Bleiberg, H., Blijham, G.H., Neijt, J.P., Veenhof, C.H., Duez, N., and Sahmoud, T. (1993). Phase II trial of mitomycin C (MMC) in advanced gallbladder and biliary tree carcinoma. An EORTC Gastrointestinal Tract Cancer Cooperative Group Study. *Ann Oncol* 4, 607-609.
- Taft, R.J., Glazov, E.A., Cloonan, N., Simons, C., Stephen, S., Faulkner, G.J., Lassmann, T., Forrest, A.R., Grimmond, S.M., Schroder, K., Irvine, K., Arakawa, T., Nakamura, M., Kubosaki, A., Hayashida, K., Kawazu, C.,

- Murata, M., Nishiyori, H., Fukuda, S., Kawai, J., Daub, C.O., Hume, D.A., Suzuki, H., Orlando, V., Carninci, P., Hayashizaki, Y., and Mattick, J.S. (2009a). Tiny RNAs associated with transcription start sites in animals. *Nat Genet* 41, 572-578.
- Taft, R.J., Glazov, E.A., Lassmann, T., Hayashizaki, Y., Carninci, P., and Mattick, J.S. (2009b). Small RNAs derived from snoRNAs. *RNA* 15, 1233-1240.
- Taipale, J., and Beachy, P.A. (2001). The Hedgehog and Wnt signalling pathways in cancer. *Nature* 411, 349-354.
- Takahashi, K., Yan, I.K., Kogure, T., Haga, H., and Patel, T. (2014). Extracellular vesicle-mediated transfer of long non-coding RNA ROR modulates chemosensitivity in human hepatocellular cancer. *FEBS Open Bio* 4, 458-467.
- Tan, S., Shi, H., Ba, M., Lin, S., Tang, H., Zeng, X., and Zhang, X. (2016). miR-409-3p sensitizes colon cancer cells to oxaliplatin by inhibiting Beclin-1-mediated autophagy. *Int J Mol Med* 37, 1030-1038.
- Tay, Y., Zhang, J., Thomson, A.M., Lim, B., and Rigoutsos, I. (2008). MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation. *Nature* 455, 1124-1128.
- Tong, Y.S., Zhou, X.L., Wang, X.W., Wu, Q.Q., Yang, T.X., Lv, J., Yang, J.S., Zhu, B., and Cao, X.F. (2014). Association of decreased expression of long non-coding RNA LOC285194 with chemoradiotherapy resistance and poor prognosis in esophageal squamous cell carcinoma. *J Transl Med* 12, 233.
- Torre, L.A., Bray, F., Siegel, R.L., Ferlay, J., Lortet-Tieulent, J., and Jemal, A. (2015). Global cancer statistics, 2012. *CA Cancer J Clin* 65, 87-108.
- Tsang, W.P., and Kwok, T.T. (2007). Riboregulator H19 induction of MDR1-associated drug resistance in human hepatocellular carcinoma cells. *Oncogene* 26, 4877-4881.
- Tsuchiya, Y., Nakajima, M., Takagi, S., Taniya, T., and Yokoi, T. (2006). MicroRNA regulates the expression of human cytochrome P4501B1. *Cancer Research* 66, 9090-9098.
- Tu, M.J., Pan, Y.Z., Qiu, J.X., Kim, E.J.H., and Yu, A.M. (2016). Impact and mechanistic role of MicroRNA-1291 on pancreatic tumorigenesis. *Journal of Clinical Oncology* 34.
- Valeri, N., Gasparini, P., Braconi, C., Paone, A., Lovat, F., Fabbri, M., Sumani, K.M., Alder, H., Amadori, D., Patel, T., Nuovo, G.J., Fishel, R., and Croce, C.M. (2010a). MicroRNA-21 induces resistance to 5-fluorouracil by down-regulating human DNA MutS homolog 2 (hMSH2). *Proc Natl Acad Sci U S A* 107, 21098-21103.
- Valeri, N., Gasparini, P., Fabbri, M., Braconi, C., Veronese, A., Lovat, F., Adair, B., Vannini, I., Fanini, F., Bottoni, A., Costinean, S., Sandhu, S.K., Nuovo, G.J., Alder, H., Gafa, R., Calore, F., Ferracin, M., Lanza, G., Volinia, S., Negrini, M., McIlhatton, M.A., Amadori, D., Fishel, R., and Croce, C.M. (2010b). Modulation of mismatch repair and genomic stability by miR-155. *Proc Natl Acad Sci U S A* 107, 6982-6987.
- Valeri, N., Vannini, I., Fanini, F., Calore, F., Adair, B., and Fabbri, M. (2009). Epigenetics, miRNAs, and human cancer: a new chapter in human gene regulation. *Mamm Genome* 20, 573-580.
- Van Der Ree, M.H., De Vree, J.M., Stelma, F., Willemse, S., Van Der Valk, M., Rietdijk, S., Molenkamp, R., Schinkel, J., Van Nuenen, A.C., Beuers, U., Hadi,

- 1936 S., Harbers, M., Van Der Veer, E., Liu, K., Grundy, J., Patick, A.K., Pavlicek,
1937 A., Blem, J., Huang, M., Grint, P., Neben, S., Gibson, N.W., Kootstra, N.A.,
1938 and Reesink, H.W. (2017). Safety, tolerability, and antiviral effect of RG-101 in
1939 patients with chronic hepatitis C: a phase 1B, double-blind, randomised
1940 controlled trial. *Lancet* 389, 709-717.
- 1941 Van Zandwijk, N., Pavlakis, N., Kao, S.C., Linton, A., Boyer, M.J., Clarke, S., Huynh,
1942 Y., Chrzanowska, A., Fulham, M.J., Bailey, D.L., Cooper, W.A., Kritharides,
1943 L., Ridley, L., Pattison, S.T., Macdiarmid, J., Brahmbhatt, H., and Reid, G.
1944 (2017). Safety and activity of microRNA-loaded minicells in patients with
1945 recurrent malignant pleural mesothelioma: a first-in-man, phase 1, open-label,
1946 dose-escalation study. *Lancet Oncol* 18, 1386-1396.
- 1947 Vishnubalaji, R., Hamam, R., Yue, S., Al-Obeed, O., Kassem, M., Liu, F.F.,
1948 Aldahmash, A., and Alajez, N.M. (2016). MicroRNA-320 suppresses
1949 colorectal cancer by targeting SOX4, FOXM1, and FOXQ1. *Oncotarget* 7,
1950 35789-35802.
- 1951 Von Schack, D., Agostino, M.J., Murray, B.S., Li, Y., Reddy, P.S., Chen, J., Choe,
1952 S.E., Strassle, B.W., Li, C., Bates, B., Zhang, L., Hu, H., Kotnis, S., Bingham,
1953 B., Liu, W., Whiteside, G.T., Samad, T.A., Kennedy, J.D., and Ajit, S.K.
1954 (2011). Dynamic changes in the microRNA expression profile reveal multiple
1955 regulatory mechanisms in the spinal nerve ligation model of neuropathic pain.
1956 *PLoS One* 6, e17670.
- 1957 Waldman, T., Kinzler, K.W., and Vogelstein, B. (1995). P21 Is Necessary for the
1958 P53-Mediated G(1) Arrest in Human Cancer-Cells. *Cancer Research* 55,
1959 5187-5190.
- 1960 Walker, T.L., White, J.D., Esdale, W.J., Burton, M.A., and Decruz, E.E. (1996).
1961 Tumour cells surviving in vivo cisplatin chemotherapy display elevated c-myc
1962 expression. *Br J Cancer* 73, 610-614.
- 1963 Wan, L.Y., Deng, J., Xiang, X.J., Zhang, L., Yu, F., Chen, J., Sun, Z., Feng, M., and
1964 Xiong, J.P. (2015). miR-320 enhances the sensitivity of human colon cancer
1965 cells to chemoradiotherapy in vitro by targeting FOXM1. *Biochem Biophys
1966 Res Commun* 457, 125-132.
- 1967 Wang, B.D., Kline, C.L., Pastor, D.M., Olson, T.L., Frank, B., Luu, T., Sharma, A.K.,
1968 Robertson, G., Weirauch, M.T., Patierno, S.R., Stuart, J.M., Irby, R.B., and
1969 Lee, N.H. (2010a). Prostate apoptosis response protein 4 sensitizes human
1970 colon cancer cells to chemotherapeutic 5-FU through mediation of an NF
1971 kappaB and microRNA network. *Mol Cancer* 9, 98.
- 1972 Wang, C.J., Stratmann, J., Zhou, Z.G., and Sun, X.F. (2010b). Suppression of
1973 microRNA-31 increases sensitivity to 5-FU at an early stage, and affects cell
1974 migration and invasion in HCT-116 colon cancer cells. *BMC Cancer* 10, 616.
- 1975 Wang, F., Li, T., Zhang, B., Li, H., Wu, Q., Yang, L., Nie, Y., Wu, K., Shi, Y., and
1976 Fan, D. (2013a). MicroRNA-19a/b regulates multidrug resistance in human
1977 gastric cancer cells by targeting PTEN. *Biochem Biophys Res Commun* 434,
1978 688-694.
- 1979 Wang, F., Li, X., Xie, X., Zhao, L., and Chen, W. (2008). UCA1, a non-protein-coding
1980 RNA up-regulated in bladder carcinoma and embryo, influencing cell growth
1981 and promoting invasion. *FEBS Lett* 582, 1919-1927.
- 1982 Wang, F., Ying, H.Q., He, B.S., Pan, Y.Q., Deng, Q.W., Sun, H.L., Chen, J., Liu, X.,
1983 and Wang, S.K. (2015a). Upregulated lncRNA-UCA1 contributes to

- 1984 progression of hepatocellular carcinoma through inhibition of miR-216b and
 1985 activation of FGFR1/ERK signaling pathway. *Oncotarget* 6, 7899-7917.
- 1986 Wang, H., Guan, Z., He, K., Qian, J., Cao, J., and Teng, L. (2017). LncRNA UCA1 in
 1987 anti-cancer drug resistance. *Oncotarget* 8, 64638-64650.
- 1988 Wang, K.C., Yang, Y.W., Liu, B., Sanyal, A., Corces-Zimmerman, R., Chen, Y.,
 1989 Lajoie, B.R., Protacio, A., Flynn, R.A., Gupta, R.A., Wysocka, J., Lei, M.,
 1990 Dekker, J., Helms, J.A., and Chang, H.Y. (2011). A long noncoding RNA
 1991 maintains active chromatin to coordinate homeotic gene expression. *Nature*
 1992 472, 120-124.
- 1993 Wang, L., Wang, Y.M., Xu, S., Wang, W.G., Chen, Y., Mao, J.Y., and Tian, B.L.
 1994 (2015b). MicroRNA-215 is upregulated by treatment with Adriamycin and
 1995 leads to the chemoresistance of hepatocellular carcinoma cells and tissues.
 1996 *Molecular Medicine Reports* 12, 5274-5280.
- 1997 Wang, L., Zhao, Z.X., Feng, W.D., Ye, Z.J., Dai, W.G., Zhang, C.H., Peng, J.J., and
 1998 Wu, K.M. (2016a). Long non-coding RNA TUG1 promotes colorectal cancer
 1999 metastasis via EMT pathway. *Oncotarget* 7, 51713-51719.
- 2000 Wang, X., Li, M., Wang, Z., Han, S., Tang, X., Ge, Y., Zhou, L., Zhou, C., Yuan, Q.,
 2001 and Yang, M. (2015c). Silencing of long noncoding RNA MALAT1 by miR-101
 2002 and miR-217 inhibits proliferation, migration, and invasion of esophageal
 2003 squamous cell carcinoma cells. *J Biol Chem* 290, 3925-3935.
- 2004 Wang, Y., Xu, Z., Jiang, J., Xu, C., Kang, J., Xiao, L., Wu, M., Xiong, J., Guo, X., and
 2005 Liu, H. (2013b). Endogenous miRNA sponge lincRNA-RoR regulates Oct4,
 2006 Nanog, and Sox2 in human embryonic stem cell self-renewal. *Dev Cell* 25,
 2007 69-80.
- 2008 Wang, Y., Zhang, D.X., Wu, K.C., Zhao, Q.C., Nie, Y.Z., and Fan, D.M. (2014). Long
 2009 Noncoding RNA MRUL Promotes ABCB1 Expression in Multidrug-Resistant
 2010 Gastric Cancer Cell Sublines. *Molecular and Cellular Biology* 34, 3182-3193.
- 2011 Wang, Y., Zhao, Y., Herbst, A., Kalinski, T., Qin, J., Wang, X., Jiang, Z., Benedix, F.,
 2012 Franke, S., Wartman, T., Camaj, P., Halangk, W., Kolligs, F.T., Jauch, K.W.,
 2013 Nelson, P.J., and Bruns, C.J. (2016b). miR-221 Mediates Chemoresistance of
 2014 Esophageal Adenocarcinoma by Direct Targeting of DKK2 Expression. *Ann*
 2015 *Surg* 264, 804-814.
- 2016 Wang, Z., Liu, M., Zhu, H., Zhang, W., He, S., Hu, C., Quan, L., Bai, J., and Xu, N.
 2017 (2013c). miR-106a is frequently upregulated in gastric cancer and inhibits the
 2018 extrinsic apoptotic pathway by targeting FAS. *Mol Carcinog* 52, 634-646.
- 2019 Weber, J., and McCormack, P.L. (2008). Panitumumab: in metastatic colorectal
 2020 cancer with wild-type KRAS. *BioDrugs* 22, 403-411.
- 2021 Wellner, U., Schubert, J., Burk, U.C., Schmalhofer, O., Zhu, F., Sonntag, A.,
 2022 Waldvogel, B., Vannier, C., Darling, D., Zur Hausen, A., Brunton, V.G.,
 2023 Morton, J., Sansom, O., Schuler, J., Stemmler, M.P., Herzberger, C., Hopt,
 2024 U., Keck, T., Brabletz, S., and Brabletz, T. (2009). The EMT-activator ZEB1
 2025 promotes tumorigenicity by repressing stemness-inhibiting microRNAs. *Nat*
 2026 *Cell Biol* 11, 1487-1495.
- 2027 Wiemer, E.a.C., Wozniak, A., Burger, H., Smid, M., Floris, G., Nzokirantevye, A.,
 2028 Sciot, R., Sleijfer, S., and Schoffski, P. (2017). Identification of microRNA
 2029 biomarkers for response of advanced soft tissue sarcomas to eribulin:
 2030 Translational results of the EORTC 62052 trial. *Eur J Cancer* 75, 33-40.

- Wilusz, J.E., Sunwoo, H., and Spector, D.L. (2009). Long noncoding RNAs: functional surprises from the RNA world. *Genes & Development* 23, 1494-1504.
- Winter, J., Jung, S., Keller, S., Gregory, R.I., and Diederichs, S. (2009). Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol* 11, 228-234.
- Wouters, M.D., Van Gent, D.C., Hoeijmakers, J.H.J., and Pothof, J. (2011). MicroRNAs, the DNA damage response and cancer. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis* 717, 54-66.
- Wu, B.Q., Jiang, Y., Zhu, F., Sun, D.L., and He, X.Z. (2017). Long Noncoding RNA PVT1 Promotes EMT and Cell Proliferation and Migration Through Downregulating p21 in Pancreatic Cancer Cells. *Technol Cancer Res Treat*, 1533034617700559.
- Wu, K., Yang, Y., Zhao, J., and Zhao, S. (2016a). BAG3-mediated miRNA let-7g and let-7i inhibit proliferation and enhance apoptosis of human esophageal carcinoma cells by targeting the drug transporter ABCC10. *Cancer Letters* 371, 125-133.
- Wu, M., Lin, Z., Li, X., Xin, X., An, J., Zheng, Q., Yang, Y., and Lu, D. (2016b). HULC cooperates with MALAT1 to aggravate liver cancer stem cells growth through telomere repeat-binding factor 2. *Sci Rep* 6, 36045.
- Xia, H., and Hui, K.M. (2014). Mechanism of cancer drug resistance and the involvement of noncoding RNAs. *Curr Med Chem* 21, 3029-3041.
- Xia, L., Zhang, D., Du, R., Pan, Y., Zhao, L., Sun, S., Hong, L., Liu, J., and Fan, D. (2008). miR-15b and miR-16 modulate multidrug resistance by targeting BCL2 in human gastric cancer cells. *Int J Cancer* 123, 372-379.
- Xia, P., and Xu, X.Y. (2015). PI3K/Akt/mTOR signaling pathway in cancer stem cells: from basic research to clinical application. *Am J Cancer Res* 5, 1602-1609.
- Xiao, B., Guo, J., Miao, Y., Jiang, Z., Huan, R., Zhang, Y., Li, D., and Zhong, J. (2009). Detection of miR-106a in gastric carcinoma and its clinical significance. *Clin Chim Acta* 400, 97-102.
- Xing, J., Wan, S., Zhou, F., Qu, F., Li, B., Myers, R.E., Fu, X., Palazzo, J.P., He, X., Chen, Z., and Yang, H. (2012). Genetic polymorphisms in pre-microRNA genes as prognostic markers of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 21, 217-227.
- Xiong, H., Ni, Z., He, J., Jiang, S., Li, X., He, J., Gong, W., Zheng, L., Chen, S., Li, B., Zhang, N., Lyu, X., Huang, G., Chen, B., Zhang, Y., and He, F. (2017). LncRNA HULC triggers autophagy via stabilizing Sirt1 and attenuates the chemosensitivity of HCC cells. *Oncogene* 36, 3528-3540.
- Xu, K., Chen, G., Qiu, Y., Yuan, Z., Li, H., Yuan, X., Sun, J., Xu, J., Liang, X., and Yin, P. (2017a). miR-503-5p confers drug resistance by targeting PUMA in colorectal carcinoma. *Oncotarget* 8, 21719-21732.
- Xu, K., Liang, X., Cui, D., Wu, Y., Shi, W., and Liu, J. (2013). miR-1915 inhibits Bcl-2 to modulate multidrug resistance by increasing drug-sensitivity in human colorectal carcinoma cells. *Mol Carcinog* 52, 70-78.
- Xu, K., Liang, X., Shen, K., Cui, D.L., Zheng, Y.H., Xu, J.H., Fan, Z.Z., Qiu, Y.Y., Li, Q., Ni, L., and Liu, J.W. (2012). miR-297 modulates multidrug resistance in human colorectal carcinoma by down-regulating MRP-2. *Biochemical Journal* 446, 291-300.

- Xu, L., Beckebaum, S., Iacob, S., Wu, G., Kaiser, G.M., Radtke, A., Liu, C., Kabar, I., Schmidt, H.H., Zhang, X., Lu, M., and Cicinnati, V.R. (2014a). MicroRNA-101 inhibits human hepatocellular carcinoma progression through EZH2 downregulation and increased cytostatic drug sensitivity. *J Hepatol* 60, 590-598.
- Xu, S., Tao, Z., Hai, B., Liang, H., Shi, Y., Wang, T., Song, W., Chen, Y., Ouyang, J., Chen, J., Kong, F., Dong, Y., Jiang, S.W., Li, W., Wang, P., Yuan, Z., Wan, X., Wang, C., Li, W., Zhang, X., and Chen, K. (2016). miR-424(322) reverses chemoresistance via T-cell immune response activation by blocking the PD-L1 immune checkpoint. *Nat Commun* 7, 11406.
- Xu, W.H., Zhang, J.B., Dang, Z., Li, X., Zhou, T., Liu, J., Wang, D.S., Song, W.J., and Dou, K.F. (2014b). Long non-coding RNA URHC regulates cell proliferation and apoptosis via ZAK through the ERK/MAPK signaling pathway in hepatocellular carcinoma. *Int J Biol Sci* 10, 664-676.
- Xu, Y., Jiang, X.M., and Cui, Y.F. (2017b). Upregulated long noncoding RNA PANDAR predicts an unfavorable prognosis and promotes tumorigenesis in cholangiocarcinoma. *Oncotargets and Therapy* 10, 2873-2883.
- Xu, Y., Leng, K.M., Li, Z.L., Zhang, F.M., Zhong, X.Y., Kang, P.C., Jiang, X.M., and Cui, Y.F. (2017c). The prognostic potential and carcinogenesis of long non-coding RNA TUG1 in human cholangiocarcinoma. *Oncotarget* 8, 65823-65835.
- Xu, Y., Xia, F., Ma, L., Shan, J., Shen, J., Yang, Z., Liu, J., Cui, Y., Bian, X., Bie, P., and Qian, C. (2011). MicroRNA-122 sensitizes HCC cancer cells to adriamycin and vincristine through modulating expression of MDR and inducing cell cycle arrest. *Cancer Lett* 310, 160-169.
- Xu, Y.J., and Fan, Y. (2015). MiR-215/192 participates in gastric cancer progression. *Clin Transl Oncol* 17, 34-40.
- Yan, H.J., Liu, W.S., Sun, W.H., Wu, J., Ji, M., Wang, Q., Zheng, X., Jiang, J.T., and Wu, C.P. (2012). miR-17-5p inhibitor enhances chemosensitivity to gemcitabine via upregulating Bim expression in pancreatic cancer cells. *Dig Dis Sci* 57, 3160-3167.
- Yan, J., Dang, Y., Liu, S., Zhang, Y., and Zhang, G. (2016). LncRNA HOTAIR promotes cisplatin resistance in gastric cancer by targeting miR-126 to activate the PI3K/AKT/MRP1 genes. *Tumour Biol*.
- Yang, G., Jiang, O., Ling, D.Q., Jiang, X.Y., Yuan, P.Z., Zeng, G., Zhu, J., Tian, J., Weng, Y.G., and Wu, D.Q. (2015). MicroRNA-522 reverses drug resistance of doxorubicin-induced HT29 colon cancer cell by targeting ABCB5. *Molecular Medicine Reports* 12, 3930-3936.
- Yang, Q., Zhang, R., Wang, X.W., Linke, S.P., Sengupta, S., Hickson, I.D., Pedrazzi, G., Perrera, C., Stagljar, I., Littman, S.J., Modrich, P., and Harris, C.C. (2004). The mismatch DNA repair heterodimer, hMSH2/6, regulates BLM helicase. *Oncogene* 23, 3749-3756.
- Yang, S.M., Huang, C., Li, X.F., Yu, M.Z., He, Y., and Li, J. (2013a). miR-21 confers cisplatin resistance in gastric cancer cells by regulating PTEN. *Toxicology* 306, 162-168.
- Yang, T., Zheng, Z.M., Li, X.N., Li, Z.F., Wang, Y., Geng, Y.F., Bai, L., and Zhang, X.B. (2013b). MiR-223 modulates multidrug resistance via downregulation of

- 2126 ABCB1 in hepatocellular carcinoma cells. *Experimental Biology and Medicine*
 2127 238, 1024-1032.
- 2128 Yang, Z., Zhou, L., Wu, L.M., Lai, M.C., Xie, H.Y., Zhang, F., and Zheng, S.S.
 2129 (2011). Overexpression of long non-coding RNA HOTAIR predicts tumor
 2130 recurrence in hepatocellular carcinoma patients following liver transplantation.
 2131 *Ann Surg Oncol* 18, 1243-1250.
- 2132 Ying, L., Chen, Q., Wang, Y., Zhou, Z., Huang, Y., and Qiu, F. (2012). Upregulated
 2133 MALAT-1 contributes to bladder cancer cell migration by inducing epithelial-
 2134 to-mesenchymal transition. *Mol Biosyst* 8, 2289-2294.
- 2135 Yiren, H., Yingcong, Y., Sunwu, Y., Keqin, L., Xiaochun, T., Senrui, C., Ende, C.,
 2136 Xizhou, L., and Yanfan, C. (2017). Long noncoding RNA MALAT1 regulates
 2137 autophagy associated chemoresistance via miR-23b-3p sequestration in
 2138 gastric cancer. *Mol Cancer* 16, 174.
- 2139 Yoon, J.H., Abdelmohsen, K., and Gorospe, M. (2014). Functional interactions
 2140 among microRNAs and long noncoding RNAs. *Seminars in Cell &*
 2141 *Developmental Biology* 34, 9-14.
- 2142 Yu, J., Ohuchida, K., Mizumoto, K., Sato, N., Kayashima, T., Fujita, H., Nakata, K.,
 2143 and Tanaka, M. (2010). MicroRNA, hsa-miR-200c, is an independent
 2144 prognostic factor in pancreatic cancer and its upregulation inhibits pancreatic
 2145 cancer invasion but increases cell proliferation. *Mol Cancer* 9, 169.
- 2146 Yu, Y., Sarkar, F.H., and Majumdar, A.P. (2013). Down-regulation of miR-21 Induces
 2147 Differentiation of Chemoresistant Colon Cancer Cells and Enhances
 2148 Susceptibility to Therapeutic Regimens. *Transl Oncol* 6, 180-186.
- 2149 Yuan, J.H., Yang, F., Wang, F., Ma, J.Z., Guo, Y.J., Tao, Q.F., Liu, F., Pan, W.,
 2150 Wang, T.T., Zhou, C.C., Wang, S.B., Wang, Y.Z., Yang, Y., Yang, N., Zhou,
 2151 W.P., Yang, G.S., and Sun, S.H. (2014). A long noncoding RNA activated by
 2152 TGF-beta promotes the invasion-metastasis cascade in hepatocellular
 2153 carcinoma. *Cancer Cell* 25, 666-681.
- 2154 Zahreddine, H., and Borden, K.L. (2013). Mechanisms and insights into drug
 2155 resistance in cancer. *Front Pharmacol* 4, 28.
- 2156 Zeng, Y., Wang, T., Liu, Y., Su, Z., Lu, P., Chen, X., and Hu, D. (2017). LncRNA
 2157 PVT1 as an effective biomarker for cancer diagnosis and detection based on
 2158 transcriptome data and meta-analysis. *Oncotarget* 8, 75455-75466.
- 2159 Zhang, B.G., Li, J.F., Yu, B.Q., Zhu, Z.G., Liu, B.Y., and Yan, M. (2012). microRNA-
 2160 21 promotes tumor proliferation and invasion in gastric cancer by targeting
 2161 PTEN. *Oncol Rep* 27, 1019-1026.
- 2162 Zhang, D., Xiao, Y.F., Zhang, J.W., Xie, R., Hu, C.J., Tang, B., Wang, S.M., Wu,
 2163 Y.Y., Hao, N.B., and Yang, S.M. (2015a). miR-1182 attenuates gastric cancer
 2164 proliferation and metastasis by targeting the open reading frame of hTERT.
 2165 *Cancer Lett* 360, 151-159.
- 2166 Zhang, E., He, X., Yin, D., Han, L., Qiu, M., Xu, T., Xia, R., Xu, L., Yin, R., and De,
 2167 W. (2016a). Increased expression of long noncoding RNA TUG1 predicts a
 2168 poor prognosis of gastric cancer and regulates cell proliferation by
 2169 epigenetically silencing of p57. *Cell Death & Disease* 7.
- 2170 Zhang, J.X., Xu, Y., Gao, Y., Chen, C., Zheng, Z.S., Yun, M., Weng, H.W., Xie, D.,
 2171 and Ye, S. (2017). Decreased expression of miR-939 contributes to
 2172 chemoresistance and metastasis of gastric cancer via dysregulation of
 2173 SLC34A2 and Raf/MEK/ERK pathway. *Molecular Cancer* 16.

- 2174 Zhang, W., Winder, T., Ning, Y., Pohl, A., Yang, D., Kahn, M., Lurje, G., Labonte,
2175 M.J., Wilson, P.M., Gordon, M.A., Hu-Lieskovan, S., Mauro, D.J., Langer, C.,
2176 Rowinsky, E.K., and Lenz, H.J. (2011). A let-7 microRNA-binding site
2177 polymorphism in 3'-untranslated region of KRAS gene predicts response in
2178 wild-type KRAS patients with metastatic colorectal cancer treated with
2179 cetuximab monotherapy. *Ann Oncol* 22, 104-109.
- 2180 Zhang, X.J., Ye, H., Zeng, C.W., He, B., Zhang, H., and Chen, Y.Q. (2010).
2181 Dysregulation of miR-15a and miR-214 in human pancreatic cancer. *J*
2182 *Hematol Oncol* 3, 46.
- 2183 Zhang, X.W., Bu, P., Liu, L., Zhang, X.Z., and Li, J. (2015b). Overexpression of long
2184 non-coding RNA PVT1 in gastric cancer cells promotes the development of
2185 multidrug resistance. *Biochemical and Biophysical Research Communications*
2186 462, 227-232.
- 2187 Zhang, X.W., Bu, P., Liu, L., Zhang, X.Z., and Li, J. (2015c). Overexpression of long
2188 non-coding RNA PVT1 in gastric cancer cells promotes the development of
2189 multidrug resistance. *Biochem Biophys Res Commun* 462, 227-232.
- 2190 Zhang, Y., Qu, X., Li, C., Fan, Y., Che, X., Wang, X., Cai, Y., Hu, X., and Liu, Y.
2191 (2015d). miR-103/107 modulates multidrug resistance in human gastric
2192 carcinoma by downregulating Cav-1. *Tumour Biol* 36, 2277-2285.
- 2193 Zhang, Y., Song, X., Wang, X., Hu, J., and Jiang, L. (2016b). Silencing of LncRNA
2194 HULC Enhances Chemotherapy Induced Apoptosis in Human Gastric Cancer.
2195 *J Med Biochem* 35, 137-143.
- 2196 Zhao, H., Wang, Y., Yang, L., Jiang, R., and Li, W. (2014a). MiR-25 promotes gastric
2197 cancer cells growth and motility by targeting RECK. *Mol Cell Biochem* 385,
2198 207-213.
- 2199 Zhao, N., Wang, R., Zhou, L., Zhu, Y., Gong, J., and Zhuang, S.M. (2014b).
2200 MicroRNA-26b suppresses the NF-kappaB signaling and enhances the
2201 chemosensitivity of hepatocellular carcinoma cells by targeting TAK1 and
2202 TAB3. *Mol Cancer* 13, 35.
- 2203 Zhao, Y., Guo, Q., Chen, J., Hu, J., Wang, S., and Sun, Y. (2014c). Role of long non-
2204 coding RNA HULC in cell proliferation, apoptosis and tumor metastasis of
2205 gastric cancer: a clinical and in vitro investigation. *Oncol Rep* 31, 358-364.
- 2206 Zheng, K., Cubero, F.J., and Nevzorova, Y.A. (2017). c-MYC-Making Liver Sick:
2207 Role of c-MYC in Hepatic Cell Function, Homeostasis and Disease. *Genes*
2208 *(Basel)* 8.
- 2209 Zheng, X., Hu, H., and Li, S. (2016). High expression of lncRNA PVT1 promotes
2210 invasion by inducing epithelial-to-mesenchymal transition in esophageal
2211 cancer. *Oncol Lett* 12, 2357-2362.
- 2212 Zhou, D.D., Liu, X.F., Lu, C.W., Pant, O.P., and Liu, X.D. (2017). Long non-coding
2213 RNA PVT1: Emerging biomarker in digestive system cancer. *Cell Prolif* 50.
- 2214 Zhou, J., Wen, Q., Li, S.F., Zhang, Y.F., Gao, N., Tian, X., Fang, Y., Gao, J., Cui,
2215 M.Z., He, X.P., Jia, L.J., Jin, H., and Qiao, H.L. (2016). Significant change of
2216 cytochrome P450s activities in patients with hepatocellular carcinoma.
2217 *Oncotarget* 7, 50612-50623.
- 2218 Zhou, X., Jin, W., Jia, H., Yan, J., and Zhang, G. (2015). MiR-223 promotes the
2219 cisplatin resistance of human gastric cancer cells via regulating cell cycle by
2220 targeting FBXW7. *J Exp Clin Cancer Res* 34, 28.

- 2221 Zhou, Y., Wan, G., Spizzo, R., Ivan, C., Mathur, R., Hu, X., Ye, X., Lu, J., Fan, F.,
2222 Xia, L., Calin, G.A., Ellis, L.M., and Lu, X. (2014). miR-203 induces oxaliplatin
2223 resistance in colorectal cancer cells by negatively regulating ATM kinase. *Mol*
2224 *Oncol* 8, 83-92.
- 2225 Zhu, Y., Luo, M., Brooks, M., Clouthier, S.G., and Wicha, M.S. (2014). Biological and
2226 clinical significance of cancer stem cell plasticity. *Clin Transl Med* 3, 32.

2227

In review




Fig.1




Fig.2

Figure 3.TIF




Fig.3




Fig.4




Fig.5

Figure 6.TIF




Fig.6




Fig.7